

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 313/06, A61K 31/365		A1	(11) International Publication Number: WO 97/16438
			(43) International Publication Date: 9 May 1997 (09.05.97)
(21) International Application Number: PCT/US96/17481		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 28 October 1996 (28.10.96)			
(30) Priority Data: 60/008,191 31 October 1995 (31.10.95) US 60/007,101 31 October 1995 (31.10.95) US 9603903.7 23 February 1996 (23.02.96) GB 9605161.0 12 March 1996 (12.03.96) GB			
(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).		<p>Published</p> <p><i>With international search report.</i></p> <p><i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
(72) Inventors; and			
(75) Inventors/Applicants (for US only): BAKER, Robert, K. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). BAO, Jianming [CN/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). KAYSER, Frank [DE/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). PARSONS, William, H. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). RUPPRECHT, Kathleen, M. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).			
(74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).			

(54) Title: TRITERPENE DERIVATIVES WITH IMMUNOSUPPRESSANT ACTIVITY



(57) Abstract

The compounds of formula (I) are useful as immunosuppressive agents.

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

TITLE OF THE INVENTION**TRITERPENE DERIVATIVES WITH IMMUNOSUPPRESSANT
ACTIVITY****5 BACKGROUND OF THE INVENTION**

Immunoregulatory abnormalities have been shown to exist in a wide variety of "autoimmune" and chronic inflammatory diseases, including systemic lupus erythematosus, chronic rheumatoid arthritis, type I and II diabetes mellitus, inflammatory bowel disease, biliary
10 cirrhosis, uveitis, multiple sclerosis and other disorders such as Crohn's disease, ulcerative colitis, bullous pemphigoid, sarcoidosis, psoriasis, ichthyosis, Graves ophthalmopathy and asthma.

Although the underlying pathogenesis of each of these conditions may be quite different, they have in common the appearance
15 of a variety of autoantibodies and self-reactive lymphocytes. Such self-reactivity may be due, in part, to a loss of the homeostatic controls under which the normal immune system operates.

Similarly, following a bone-marrow or an organ transplantation, the host lymphocytes recognize the foreign tissue
20 antigens and begin to produce antibodies which lead to graft rejection.

One end result of an autoimmune or a rejection process is tissue destruction caused by inflammatory cells and the mediators they release. Anti-inflammatory agents such as NSAID's act principally by blocking the effect or secretion of these mediators but do nothing to
25 modify the immunologic basis of the disease. On the other hand, cytotoxic agents, such as cyclophosphamide, act in such a nonspecific fashion that both the normal and autoimmune responses are shut off. Indeed, patients treated with such nonspecific immunosuppressive agents are as likely to succumb from infection as they are from their
30 autoimmune disease.

Cyclosporin A (CsA), which was approved by the US FDA in 1983 is currently the leading drug used to prevent rejection of transplanted organs. In 1993, FK-506 (Prograf) was approved by the US FDA for the prevention of rejection in liver transplantation. CsA

- 2 -


and FK-506 act by inhibiting the body's immune system from mobilizing its vast arsenal of natural protecting agents to reject the transplant's foreign protein. In 1994, CsA was approved by the US FDA for the treatment of severe psoriasis and has been approved by

5 European regulatory agencies for the treatment of atopic dermatitis. Though they are effective in fighting transplant rejection, CsA and FK-506 are known to cause several undesirable side effects including nephrotoxicity, neurotoxicity, and gastrointestinal discomfort.

Newer, safer drugs exhibiting less side effects are

10 constantly being searched for in the field.

Four active components of *Spachea correa* were recently identified which inhibit thymidine uptake of T cells and are useful as immunosuppressive agents in animals, including man.



Formula 1(a) b is a single bond and R is OAc

Formula 1(b) b is a double bond and R is OAc

Formula 1(c) b is a single bond and R is OH

Formula 1(d) b is a double bond and R is OH

15


These compounds are useful as immunosuppressive agents in animals, including man. The present invention describes newly developed immunosuppressive compounds derived from the compounds described in Formulae 1(a) through 1(d) and which have the relative

20 stereochemistry depicted above.

- 3 -

SUMMARY OF THE INVENTION

A class of triterpene derivatives of the general structural Formula I



I

are useful as immunosuppressants.

As immunosuppressants, the compounds of this invention are useful in the treatment of autoimmune diseases, the prevention of rejection of foreign organ transplants and/or related afflictions, diseases and illnesses. Also within the scope of this invention are pharmaceutical formulations comprising a compound of Formula I and a pharmaceutical carrier, as well as, pharmaceutical formulations comprising a compound of Formula I, a second immunosuppressant compound and a pharmaceutical carrier.

DETAILED DESCRIPTION OF THE INVENTIONA. Scope of the Invention

The present invention is related to compounds of formula I, including but not limited to those specified in the examples, which are useful in a mammalian subject for the treatment and prevention of the resistance by transplantation of organs or tissue, graft-versus-host diseases brought about by medulla ossium transplantation; rheumatoid arthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, type I diabetes uveitis, juvenile-onset or recent-onset diabetes mellitus, posterior uveitis, allergic

- 4 -


encephalomyelitis, glomerulonephritis, infectious diseases caused by pathogenic microorganisms, inflammatory and hyperproliferative skin diseases, psoriasis, atopic dermatitis, contact dermatitis, eczematous dermatitis, seborrhoeic dermatitis, Lichen planus, Pemphigus, bullous pemphigoid, Epidermolysis bullosa, urticaria, angioedemas, vasculitides, erythemas, cutaneous eosinophilias, Lupus erythematosus, acne, Alopecia areata, keratoconjunctivitis, vernal conjunctivitis, uveitis associated with Behcet's disease, keratitis, herpetic keratitis, conical cornea, dystrophia epithelialis corneae, corneal leukoma, ocular pemphigus, Mooren's ulcer, Scleritis, Graves' ophthalmopathy, Vogt-Koyanagi-Harada syndrome, sarcoidosis, etc.; pollen allergies, reversible obstructive airway disease, bronchial asthma, allergic asthma, intrinsic asthma, extrinsic asthma and dust asthma, chronic or inveterate asthma, late asthma and airway hyper-responsiveness, bronchitis, gastric ulcers, vascular damage caused by ischemic diseases and thrombosis, ischemic bowel diseases, inflammatory bowel diseases, necrotizing enterocolitis, intestinal lesions associated with thermal burns and leukotriene B₄-mediated diseases, Coeliac diseases, proctitis, eosinophilic gastroenteritis, mastocytosis, Crohn's disease, ulcerative colitis, migraine, rhinitis, eczema, interstitial nephritis, Good-pasture's syndrome, hemolytic-uremic syndrome, diabetic nephropathy, multiple myositis, Guillain-Barre syndrome, Meniere's disease, polyneuritis, multiple neuritis, mononeuritis, radiculopathy, hyperthyroidism, Basedow's disease, pure red cell aplasia, aplastic anemia, hypoplastic anemia, idiopathic thrombocytopenic purpura, autoimmune hemolytic anemia, agranulocytosis, pernicious anemia, megaloblastic anemia, anerythroplasia, osteoporosis, sarcoidosis, fibroid lung, idiopathic interstitial pneumonia, dermatomyositis, leukoderma vulgaris, ichthyosis vulgaris, photoallergic sensitivity, cutaneous T cell lymphoma, arteriosclerosis, atherosclerosis, aortitis syndrome, polyarteritis nodosa, myocarditis, scleroderma, Wegener's granuloma, Sjogren's syndrome, adiposis, eosinophilic fascitis, lesions of gingiva, periodontium, alveolar bone, substantia ossea dentis, glomerulonephritis, male pattern alopecia or alopecia senilis by preventing epilation or providing hair germination

- 5 -

and/or promoting hair generation and hair growth; muscular dystrophy; Pyoderma and Sezary's syndrome, Addison's disease, ischemia-reperfusion injury of organs which occurs upon preservation, transplantation or ischemic disease, for example, thrombosis and cardiac infraction, endotoxin-shock, pseudomembranous colitis, colitis caused by drug or radiation, ischemic acute renal insufficiency, chronic renal insufficiency, toxinosis caused by lung-oxygen or drug, for example, paracort and bleomycins), lung cancer, pulmonary emphysema, cataracta, siderosis, retinitis, pigmentosa, senile macular degeneration, vitreal scarring, corneal alkali bum; dermatitis erythema multiforme, linear IgA ballous dermatitis and cement dermatitis, gingivitis, periodontitis, sepsis, pancreatitis, diseases caused by environmental pollution, aging, carcinogenesis, metastasis of carcinoma and hypobaropathy; disease caused by histamine or leukotriene-C4 release; Behcet's disease, autoimmune hepatitis, primary biliary cirrhosis sclerosing cholangitis), partial liver resection, acute liver necrosis, necrosis caused by toxin, viral hepatitis, shock, or anoxia, B-virus hepatitis, non-A/non-B hepatitis, cirrhosis, alcoholic cirrhosis, hepatic failure, fulminant hepatic failure, late-onset hepatic failure, "acute-on-chronic" liver failure, augmentation of chemotherapeutic effect, preventing or treating activity of cytomegalovirus infection, HCMV infection, and antiinflammatory activity; and treatment of immunodepression or a disorder involving immunodepression, including AIDS, cancer, senile dementia, trauma, chronic bacterial infection, and certain central nervous system disorders.

More particularly this invention relates to compounds of the general structural formula I:

- 6 -



I

or a pharmaceutically acceptable salt, crystal form or hydrate, wherein:

5 X is: O; S, NH, H and R^1 ;

a is: a single bond, or a double bond when R^4 is absent;

b and c are independently: a single bond, or a double bond;

10

n is: 1 to 4;

m is: 1 to 4;

15 r is: 0 or 1;

s is: 0 or 1;

R^1 and R^2 are independently:

20

a) H, or

b) (C_1-C_6) -alkyl, wherein alkyl is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C_1-C_6) -alkoxy, vinyl, cyano, oxo, nitro, hydroxy, CHO, CO_2H , COC_1-C_6 -alkyl, $CO_2C_1-C_6$ -alkyl, $CONR^1R^2$, NR^1R^2 , $NR^1COC_1-C_6$ -alkyl, aryl, wherein aryl is defined as phenyl or naphthyl,

25

- 7 -

5 unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, phenyl, phenoxy, cyano, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl and any two of adjacent substituents can be joined to form a 5-, 6- or 7-membered fused ring said ring containing 1 or 2 oxygen atoms and the remainder carbon atoms, heteroaryl, wherein heteroaryl is defined as a 5 or 6-membered ring substituted with one and two heteroatoms selected from O, S, N, unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, cyano, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, any two adjacent substituents can be joined to form a 5-, 6- or 7-membered fused ring said ring containing 1 or 2 oxygen atoms and the remainder carbon atoms, or any two adjacent substituents can be joined together to form a benzo-fused ring;

20

R³ is:

- a) -(C₁-C₆)-alkyl, alkyl as defined above;
- b) -(C₁-C₆)-alkenyl, wherein alkenyl is unsubstituted or substituted with one, two or three substituents selected from from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, aryl as defined above, and heteroaryl as defined above;
- c) -(C₁-C₆)-alkynyl, wherein alkynyl is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, aryl as defined above, and heteroaryl as defined above,

- 8 -


- d) -aryl, aryl as defined above, or
- e) -heteroaryl, heteroaryl as defined above;

R⁴ is:

- 5 a) absent and a is a double bond;
- b) -H,
- c) -OH,
- d) =O,
- e) -O[(C=O)O_r]_sC₁-C₁₀-alkyl, alkyl as defined above,
- 10 f) -O[(C=O)O_r]_sC₂-C₁₀-alkenyl, as defined above,
- g) -O[(C=O)O_r]_sC₂-C₆-alkynyl, alkynyl as defined above,
- h) -O[(C=O)O_r]_s(C₃-C₇)-cycloalkyl,
- i) -O[(C=O)O_r]_saryl, aryl as defined above,
- j) -O[(C=O)O_r]_sheteroaryl, heteroaryl as defined above,
- 15 k) -O(CH₂)_nO(CH₂)_mheteroaryl, heteroaryl as defined above,
- l) -O(CH₂)_nO(CH₂)_maryl, aryl as defined above,
- m) -OC(=O)NR¹R²,
- n) -OSO₂R³, or
- o) -NR¹R².

20

An embodiment of the invention are the compounds of structural Formula I,



I

- 25 or a pharmaceutically acceptable salt, crystal form or hydrate, wherein:

- 9 -

X is: O, S, or NH;

a is: a single bond;

5

b and c are independently: a single bond or a double bond;

n is: 1 to 4;

10

m is: 1 to 4;

r is: 0 or 1;

s is: 0 or 1;

15

R¹ and R² are independently:

a) H, or

20 b) (C₁-C₆)-alkyl, wherein alkyl is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, vinyl, cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, aryl, wherein aryl is defined as phenyl or naphthyl, unsubstituted or substituted with one, two or three
25 substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, phenyl, phenoxy, cyano, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl and any two of adjacent substituents can be joined to form a 5-, 6- or 7-
30 membered fused ring said ring containing 1 or 2 oxygen atoms and the remainder carbon atoms, heteroaryl, wherein heteroaryl is defined as a 5 or 6-membered ring substituted with one and two heteroatoms selected from O, S, N, unsubstituted or substituted with one, two or three

- 10 -

substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, cyano, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, any two adjacent substituents can be joined to form a 5-, 6- or 7-membered fused ring said ring containing 1 or 2 oxygen atoms and the remainder carbon atoms, or any two adjacent substituents can be joined together to form a benzo-fused ring;

10 R³ is:

- a) -(C₁-C₆)-alkyl, alkyl as defined above;
- b) -(C₁-C₆)-alkenyl, wherein alkenyl is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, aryl as defined above, and heteroaryl as defined above;
- c) -(C₁-C₆)-alkynyl, wherein alkynyl is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, aryl as defined above, and heteroaryl as defined above,
- d) -aryl, aryl as defined above, or
- e) -heteroaryl, heteroaryl as defined above;


R⁴ is:

- a) absent and a is a double bond;
- b) -H,
- c) -OH,
- d) =O,
- e) -O[(C=O)O_r]_sC₁-C₁₀-alkyl, alkyl as defined above,
- f) -O[(C=O)O_r]_sC₂-C₁₀-alkenyl, as defined above,
- g) -O[(C=O)O_r]_sC₂-C₆-alkynyl, alkynyl as defined above,

- 11 -

- h) $-O[(C=O)O_r]_s(C_3-C_7)\text{-cycloalkyl}$,
 i) $-O[(C=O)O_r]_s\text{aryl}$, aryl as defined above,
 j) $-O[(C=O)O_r]_s\text{heteroaryl}$, heteroaryl as defined above,
 k) $-O(CH_2)_nO(CH_2)_m\text{heteroaryl}$, heteroaryl as defined above,
 5 l) $-O(CH_2)_nO(CH_2)_m\text{aryl}$, aryl as defined above,
 m) $-OC(=O)NR^1R^2$,
 n) $-OSO_2R^3$, or
 o) $-NR^1R^2$.

10 An embodiment of this embodiment of the invention are the compounds of structural Formula I,



I

or a pharmaceutically acceptable salt, crystal form or hydrate, wherein:

15

X is: O;

a is: a single bond;

20 b and c are independently: a single bond or a double bond;

n is: 1 to 4;

m is: 1 to 4;

25

- 12 -

r is: 0 or 1;

s is: 0 or 1;

5 R¹ and R² are independently:

a) H, or

10 b) (C₁-C₆)-alkyl, wherein alkyl is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, vinyl, cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, aryl, wherein aryl is defined as phenyl or naphthyl, unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, 15 F, I, (C₁-C₆)-alkoxy, phenyl, phenoxy, cyano, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl and any two of adjacent substituents can be joined to form a 5-, 6- or 7-membered fused ring said ring containing 1 or 2 oxygen atoms and the remainder carbon atoms, heteroaryl, wherein 20 heteroaryl is defined as a 5 or 6-membered ring substituted with one and two heteroatoms selected from O, S, N, unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, 25 F, I, (C₁-C₆)-alkoxy, cyano, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, any two adjacent substituents can be joined to form a 5-, 6- or 7-membered fused ring said ring containing 1 or 2 oxygen atoms and the remainder carbon atoms, or any two adjacent substituents can be joined 30 together to form a benzo-fused ring;

- 13 -

R³ is:

- a) -(C₁-C₆)-alkyl, alkyl as defined above,
- b) -aryl, aryl as defined above, or
- c) -heteroaryl, heteroaryl as defined above;

5

R⁴ is:

- a) -O[(C=O)O_R]_sC₁-C₁₀-alkyl, alkyl as defined above,
- b) -O[(C=O)O_R]_s(C₃-C₇)-cycloalkyl,
- c) -O[(C=O)O_R]_saryl, aryl as defined above,
- 10 d) -O[(C=O)O_R]_sheteroaryl, heteroaryl as defined above,
- e) -O(CH₂)_nO(CH₂)_mheteroaryl, heteroaryl as defined above,
- f) -O(CH₂)_nO(CH₂)_maryl, aryl as defined above,
- g) -OC(=O)NR¹R², or
- h) -OSO₂R³.

15

A preferred embodiment of this embodiment are the compounds of structural Formula I, as recited in above, or a pharmaceutically acceptable salt, crystal form or hydrate, wherein:


20 R⁴ is:

- a) -O[(C=O)O_R]_saryl, wherein aryl is defined as phenyl or naphthyl, unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, phenyl, phenoxy, cyano, nitro, 25 hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl and any two of adjacent substituents can be joined to form a 5-, 6- or 7-membered fused ring said ring containing 1 or 2 oxygen atoms and the remainder carbon atoms, or
- 30 b) -O[(C=O)O_R]_sheteroaryl, wherein heteroaryl is defined as a 5 or 6-membered ring substituted with one and two heteroatoms selected from O, S, N, unsubstituted or substituted with one, two or three substituents selected from

- 14 -

the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, cyano, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, any two adjacent substituents can be joined to form a 5-, 6- or 7-membered fused ring said ring containing 1 or 2 oxygen atoms and the remainder carbon atoms, or any two adjacent substituents can be joined together to form a benzo-fused ring.

Another embodiment of this invention are the compounds of structural Formula I,



I

or a pharmaceutically acceptable salt, crystal form or hydrate, wherein:

X is: H and R¹;

a is: a single bond;

b and c are independently: a single bond or a double bond;

n is: 1 to 4;

m is: 1 to 4;

- 15 -

r is: 0 or 1;

s is: 0 or 1;

5 R¹ and R² are independently:

a) H, or

10 b) (C₁-C₆)-alkyl, wherein alkyl is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, vinyl, cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, aryl, wherein aryl is defined as phenyl or naphthyl, unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, 15 F, I, (C₁-C₆)-alkoxy, phenyl, phenoxy, cyano, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl and any two of adjacent substituents can be joined to form a 5-, 6- or 7-membered fused ring said ring containing 1 or 2 oxygen atoms and the remainder carbon atoms, heteroaryl, wherein heteroaryl is defined as a 5 or 6-membered ring substituted with one and two heteroatoms selected from O, S, N, unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, 25 F, I, (C₁-C₆)-alkoxy, cyano, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, any two adjacent substituents can be joined to form a 5-, 6- or 7-membered fused ring said ring containing 1 or 2 oxygen atoms and the remainder carbon atoms, or any two adjacent substituents can be joined 30 together to form a benzo-fused ring;

R³ is:


a) -(C₁-C₆)-alkyl, alkyl as defined above;

- 16 -

- 5 b) -(C₁-C₆)-alkenyl, wherein alkenyl is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, aryl as defined above, and heteroaryl as defined above;
- 10 c) -(C₁-C₆)-alkynyl, wherein alkynyl is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, aryl as defined above, and heteroaryl as defined above,
- 15 d) -aryl, aryl as defined above, or
- e) -heteroaryl, heteroaryl as defined above;
- R⁴ is:
- a) absent and a is a double bond;
- b) -H,
- c) -OH,
- 20 d) =O,
- e) -O[(C=O)O_r]_sC₁-C₁₀-alkyl, alkyl as defined above,
- f) -O[(C=O)O_r]_sC₂-C₁₀-alkenyl, as defined above,
- g) -O[(C=O)O_r]_sC₂-C₆-alkynyl, alkynyl as defined above,
- h) -O[(C=O)O_r]_s(C₃-C₇)-cycloalkyl,
- 25 i) -O[(C=O)O_r]_saryl, aryl as defined above,
- j) -O[(C=O)O_r]_sheteroaryl, heteroaryl as defined above,
- k) -O(CH₂)_nO(CH₂)_mheteroaryl, heteroaryl as defined above,
- l) -O(CH₂)_nO(CH₂)_maryl, aryl as defined above,
- m) -OC(=O)NR¹R²,
- 30 n) -OSO₂R³, or
- o) -NR¹R².

An embodiment of this embodiment are the compounds of structural Formula I,

- 17 -



I

or a pharmaceutically acceptable salt, crystal form or hydrate, wherein:

5 X is: H and R¹;

a is: a single bond;

b and c are independently: a single bond or a double bond;

10

n is: 1 to 4;

m is: 1 to 4;

15 r is: 0 or 1;

s is: 0 or 1;

R¹ and R² are independently:

20

a) H, or

b) (C₁-C₆)-alkyl, wherein alkyl is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, vinyl, cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, aryl, wherein aryl is defined as phenyl or naphthyl,

25

- 18 -

5 unsubstituted or substituted with one, two or three
substituents selected from the group consisting of: Br, Cl,
F, I, (C₁-C₆)-alkoxy, phenyl, phenoxy, cyano, nitro,
hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl,
CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl and any two of
adjacent substituents can be joined to form a 5-, 6- or 7-
membered fused ring said ring containing 1 or 2 oxygen
atoms and the remainder carbon atoms, heteroaryl, wherein
heteroaryl is defined as a 5 or 6-membered ring substituted
with one and two heteroatoms selected from O, S, N,
unsubstituted or substituted with one, two or three
substituents selected from the group consisting of: Br, Cl,
F, I, (C₁-C₆)-alkoxy, cyano, nitro, hydroxy, CHO, CO₂H,
COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R²,
NR¹COC₁-C₆-alkyl, any two adjacent substituents can be
joined to form a 5-, 6- or 7-membered fused ring said ring
containing 1 or 2 oxygen atoms and the remainder carbon
atoms, or any two adjacent substituents can be joined
together to form a benzo-fused ring;

20

R³ is:

- a) -(C₁-C₆)-alkyl, alkyl as defined above;
- b) -(C₁-C₆)-alkenyl, wherein alkenyl is unsubstituted or
substituted with one, two or three substituents selected from
from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy,
cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl,
CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl,
aryl as defined above, and heteroaryl as defined above;
- c) -(C₁-C₆)-alkynyl, wherein alkynyl is unsubstituted or
substituted with one, two or three substituents selected from
the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy,
cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl,
CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl,
aryl as defined above, and heteroaryl as defined above,

- 19 -

- d) -aryl, aryl as defined above, or
- e) -heteroaryl, heteroaryl as defined above;

R⁴ is:

- 5 a) -OH,
- b) -O[(C=O)O_r]_sC₁-C₁₀-alkyl, alkyl as defined above,
- c) -O[(C=O)O_r]_s(C₃-C₇)-cycloalkyl,
- d) -O[(C=O)O_r]_saryl, aryl as defined above,
- e) -O[(C=O)O_r]_sheteroaryl, heteroaryl as defined above,
- 10 f) -O(CH₂)_nO(CH₂)_mheteroaryl, heteroaryl as defined above,
- g) -O(CH₂)_nO(CH₂)_maryl, aryl as defined above,
- h) -OC(=O)NR¹R², or
- i) -OSO₂R³.

15 An embodiment of the invention is a compound selected from the group consisting of:

4,6,7,15,16-pentakis(acetyloxy)-18-hydroxy-22-methoxycarbonyl-
[6 α ,7 α ,15 β ,16 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-
20 1,20(29),21-trien-3-one;

4-(2-bromobenzoyl)oxy-6,7,15,16-tetrakis(acetyloxy)-18-hydroxy-22-
methoxycarbonyl[6 α ,7 α ,15 β ,16 β]D:A-Friedo-A-homo-27,30-dinor-24-
oxaoleana-1,20(29),21-trien-3-one;

25 4,6,7,15,16-pentakis(acetyloxy)-18-hydroxy-22-methoxycarbonyl-
[6 α ,7 α ,15 β ,16 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-
1,20(29),21-triene;

30 4,6,7,15,16-pentakis(acetyloxy)-21,22-epoxy-18-hydroxy-22-
methoxycarbonyl[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-
dinor-24-oxaoleana-20(29)-en-3-one;

- 20 -

4,6,7,15,16-pentakis(acetyloxy)-18-hydroxy-22-methoxycarbonyl-
[6 α ,7 α ,15 β ,16 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-
20(29),21-dien-3-one;

5 4-(2-bromobenzoyl)oxy-6,7,15,16-tetrakis(acetyloxy)-18-hydroxy-22-
methoxycarbonyl[6 α ,7 α ,15 β ,16 β]D:A-Friedo-A-homo-27,30-dinor-24-
oxaoleana-20(29),21-dien-3-one;

6,7,15,16-tetrakis(acetyloxy)-21,22-epoxy-4,18-dihydroxy-22-
10 methoxycarbonyl[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-
dinor-24-oxaoleana-20(29)-en-3-one; and

4-(2-bromobenzoyl)oxy-6,7,15,16-tetrakis(acetyloxy)-21,22-epoxy-18-
hydroxy-22-methoxycarbonyl[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-
15 homo-27,30-dinor-24-oxaoleana-20(29)-en-3-one.

The compounds of the present invention have asymmetric centers and this invention includes all of the optical isomers and mixtures thereof.

20 In addition compounds with carbon-carbon double bonds may occur in Z- and E- forms with all isomeric forms of the compounds being included in the present invention.

As used herein, the term "alkyl" includes those alkyl groups of a designated number of carbon atoms of either a straight, branched, or cyclic configuration. Examples of "alkyl" include methyl, ethyl, 25 propyl, isopropyl, butyl, sec-and tert-butyl, pentyl, hexyl, heptyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, norbornyl, and the like. "Alkoxy" represents an alkyl group of indicated number of carbon atoms attached through an oxygen bridge, 30 such as methoxy, ethoxy, propoxy, butoxy and pentoxy.

"Alkenyl" is intended to include hydrocarbon chains of a specified number of carbon atoms of either a straight- or branched-configuration and at least one unsaturation, which may occur at any point along the chain, such as ethenyl, propenyl, butenyl, pentenyl,

- 21 -

dimethyl pentenyl, and the like, and includes E and Z forms, where applicable. "Halogen", as used herein, means fluoro, chloro, bromo and iodo.

The term "aryl" is defined as a phenyl or naphthyl ring
5 which is optionally substituted at any available carbon atoms with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, phenyl, phenoxy, cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl. The aryl may also be substituted with a fused
10 5-, 6-, or 7-membered ring containing one or two oxygens and the remaining ring atoms being carbon, the fused 5-, 6-, or 7-ring being selected from the group consisting of: dioxolanyl, dihydrofuranyl, dihydropyranyl, and dioxanyl.

The term "heteroaryl" as utilized herein is intended to
15 include the following a 5 or 6-membered ring substituted with one or two heteroatoms selected from O, S, N, and is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, cyano, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R²,
20 NR¹COC₁-C₆-alkyl, any two adjacent substituents can be joined to form a 5-, 6- or 7-membered fused ring said ring containing 1 or 2 oxygen atoms and the remainder carbon atoms, or any two adjacent substituents can be joined together to form a benzo-fused ring. Heteroaryl groups within the scope of this definition include but are not limited to:
25 acridinyl, carbazolyl, cinnolinyl, quinoxalinyl, pyrazolyl, indolyl, benzotriazolyl, furanyl, thienyl, benzothienyl, benzofuranyl, quinolinyl, isoquinolinyl, pyrazinyl, pyridazinyl, pyridinyl, pyrimidinyl, and pyrrolyl which are substituted or unsubstituted as defined above.


In the compounds of Formula I, the heteroaryl group may
30 be optionally substituted with the substituents listed above at any available carbon atom or nitrogen atom (if present), but compounds bearing certain substituents, directly substituted to a nitrogen may be relatively unstable and are not preferred. The heteroaryl may also be fused to a second 5-, 6-, or 7-membered ring containing one or two

- 22 -

oxygen selected from the , the remaining ring atoms being carbon, selected from the group consisting of: dioxolanyl, dihydrofuranyl, dihydropyranyl, and dioxanyl.

Pharmaceutically acceptable salts include both the metallic
5 (inorganic) salts and organic salts; a list of which is given in
Remington's Pharmaceutical Sciences, 17th Edition, pg. 1418 (1985). It
is well known to one skilled in the art that an appropriate salt form is
chosen based on physical and chemical stability, flowability, hydro-
scopicity and solubility. As will be understood by those skilled in the
10 art, pharmaceutically acceptable salts include, but are not limited to salts
of inorganic acids such as hydrochloride, sulfate, phosphate,
diphosphate, hydrobromide, and nitrate or salts of an organic acid such
as malate, maleate, fumarate, tartrate, succinate, citrate, acetate, lactate,
methanesulfonate, p-toluenesulfonate or palmoate, salicylate and
15 stearate. Similarly pharmaceutically acceptable cations include, but are
not limited to sodium, potassium, calcium, aluminum, lithium and
ammonium (especially ammonium salts with secondary amines).
Preferred salts of this invention for the reasons cited above include
potassium, sodium, calcium and ammonium salts. Also included within
20 the scope of this invention are crystal forms, hydrates and solvates of
the compounds of Formula I.

- 23 -

REACTION SCHEME A


- 5 As seen in Scheme A, compound I, 4,5,6,15,16-pentakis
 (acetyloxy)-21,22-epoxy-18-hydroxy-22-methoxycarbonyl[6 α , 7 α ,
 15 β , 16 β , 21 β , 22 β]D:A-Freido-A-homo-27,30-dinor-24-oxaoleana-
 1,20(29)-diene-3-one, isolated from *Spachea correa* in liquid ammonia
 with lithium metal will result in the reduction of the C1 olefin group to
 10 produce the saturated lactone. Alternative methods for reducing the C1
 olefin group and/or the C20(29) olefin that are known in the art may
 also be employed. US Serial Number 08/476,806 filed on June 7, 1995
 describes the isolation of compound I and is hereby incorporated by

- 24 -

reference. The resultant lactone can then be converted to the oxepin analog by procedures described in Reaction Scheme B.

- It should also be noted that compounds of Formula I having the 11,12-double bond can be prepared using the starting material,
- 5 4,6,7,15,16-pentakisacetoxy-21,22-epoxy-18-hydroxy-22-methoxycarbonyl[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Freido-A-homo-27,30-dinor-24-oxaleana-1,11,20(29)-trien-3-one, isolated from *Spachea*
- correa and following the procedures described herein. However, there may be reactions where it will not be possible to selectively operate on
- 10 one of the double bonds, for example, ozonolysis.



REACTION SCHEME B



- 25 -


As seen in Scheme B, compound I [(4,6,7,15,16-pentakis(acetyloxy)-21,22-epoxy-18-hydroxy-22-methoxycarbonyl-[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Freido-A-homo-27,30-dinor-24-oxaoleana-1, 20(29)-dien-3-one] , isolated from *Spachea correa* can be converted to its oxepin analog in a two step process. US Serial Number 08/476,806 filed June 7, 1995, describing the isolation of compound I and is hereby incorporated by reference. Lactone I is first reduced to the lactol. This can be accomplished by using a variety of reducing agents including di-isobutylaluminum hydride (DIBAL-H) and sodium bis(2-methoxyethoxy)aluminum hydride (Red-Al). A more optimal reducing agent is the use of lithium tri-t-butoxyaluminum hydride in an inert solvent such as dichloromethane at reduced temperatures, preferably 0°C. The purified lactol intermediate is then reacted with triethylsilane and a Lewis acid such as borontrifluoride diethyl etherate to give the ether (oxepin) analog of I.

- 26 -

REACTION SCHEME C


- In a variation of Scheme B, ether (oxepin) derivatives substituted at C3 can also be prepared. Thus in Reaction Scheme C, lactone I is first reduced to the lactol as described in Reaction Scheme B. The purified lactol intermediate is then reacted with a trialkylaluminum reagent, as exemplified in this scheme by triethylaluminum (Et_3Al) to give the ethyl derivative. The allyl derivative can be prepared with allyltrimethylsilane and a Lewis acid, such as borontrifluoride diethyl etherate.

- 27 -

REACTION SCHEME D


- 5 The C21-C22 epoxide of lactone or ether derivatives can be converted to the olefin by use of a WCl_6/BuLi complex (1:2) in tetrahydrofuran (THF) by procedures developed by Sharpless *et al.* (*J. Am. Chem. Soc.*, 94, 6538-6540, 1972). This conversion can be achieved before or after any of the reaction schemes described.

- 28 -

REACTION SCHEME E

- Lactone or ether derivatives can be selectively de-acetylated
- 5 at C4 to give the corresponding alcohol by reacting it with an aqueous solution of HCl (preferably 2M to 3M concentration) in THF. It can also be prepared by reacting I with CH₃(Cl)Al[N(OCH₃)CH₃] (Weinreb reagent) in inert solvents such as THF, toluene or methylene chloride.
- If a product from this reaction contains the epoxide, it can
- 10 be removed by the method described in Reaction Scheme D.


- 29 -

REACTION SCHEME F

- The C4 hydroxy group can be oxidized to the corresponding ketone by a variety of oxidizing agents. The Jones reagent (chromic acid and sulfuric acid in H₂O), pyridinium chlorochromate, and oxalyl chloride plus DMSO all will achieve this conversion.
- 5


- 30 -

REACTION SCHEME G




5 The C4 hydroxy group can be dehydrated to give the olefin. Reaction of the alcohol with tris-phenoxyethylphosphonium iodide in hexamethylphosphorous triamide (HMPT) at 75°C will achieve this conversion.

- 31 -

REACTION SCHEME H

- 32 -

REACTION SCHEME H (CONT'D)

- As depicted in Reaction Scheme H, esters at C4 can be
- 5 prepared by reaction of a preformed carboxylic acid chloride with the C4 alcohol derivative (Reaction Scheme E) in a basic solvent such as pyridine. It should be understood that R^4 is used to represent a portion of the R^4 definition, e.g. R^4 can be an alkyl carbonate which is depicted in the scheme as $\text{OC}(=\text{O})\text{OR}^4$, R^4 representing the alkyl substituent.
 - 10 The acid chlorides, when not purchased, are prepared by stirring the carboxylic acids in reagents such as oxalyl chloride or thionyl chloride. Esters may also be prepared by reaction of the acid chloride and C4


- 33 -

alcohol with silver cyanide (AgCN) in an aprotic solvent such as HMPA. C4 sulfonate derivatives are prepared in a similar manner by reaction with sulfonyl chlorides.

- 5 C4 carbonate and carbamate derivatives are prepared by first reacting the C4 alcohol derivative with carbonyldiimidazole (CDI) to obtain the imidazolecarbonyl intermediate which is then reacted with an alcohol or amine (R^1R^2NH) to give the corresponding carbonate or carbamate derivatives.

- 10 C4 ether derivatives can also be prepared. The best procedure involves reacting an alcohol with trifluoromethanesulfonic anhydride (Tf_2O , triflic anhydride) to obtain the preformed triflate in dichloromethane at reduced temperature, preferably $-78^\circ C$. To this solution is added the triterpene alcohol, the reaction mixture is warmed to room temperature and stirring is continued until reaction is complete.
- 15 Ethers may also be prepared by heating a mixture of triterpene C4 alcohol, the appropriate alkylhalide and an excess of silver oxide (Ag_2O) in an aprotic inert solvent such as THF.

- 34 -

REACTION SCHEME I

Amines at C4 can be prepared from the C4 ketone described in Reaction Scheme F by reaction with an amine NHR^1R^2 in a variety of solvents with a reducing agent such as sodium cyanoborohydride.

UTILITY

The present invention is related to compounds of formula I, including but not limited to those specified in the examples, which are useful in a mammalian subject for the treatment and prevention of immunemediated diseases such as the resistance by transplantation of organs or tissue such as heart, kidney, liver, medulla ossium, skin, cornea, lung, pancreas, intestinum tenue, limb, muscle,

- 35 -

nervus, duodenum, small-bowel, pancreatic-islet-cell, including xeno transplants, etc.; graft-versus-host diseases brought about by medulla ossium transplantation; autoimmune diseases such as rheumatoid arthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, type I diabetes uveitis, juvenile-onset or recent-onset diabetes mellitus, posterior uveitis, allergic encephalomyelitis, glomerulonephritis, and the like; and further infectious diseases caused by pathogenic microorganisms. Further uses may include the treatment and prophylaxis of inflammatory and hyperproliferative skin diseases and cutaneous manifestations of immunologically mediated illnesses, such as psoriasis, atopic dermatitis, contact dermatitis and further eczematous dermatitises and further eczematous dermatitises, seborrhoeis dermatitis, Lichen planus, Pemphigus, bullous pemphigoid, Epidermolysis bullosa, urticaria, angioedemas, vasculitides, erythemas, cutaneous eosinophilias, Lupus erythematosus, acne and Alopecia areata; various eye diseases (autoimmune and otherwise) such as keratoconjunctivitis, vernal conjunctivitis, uveitis associated with Behcet's disease, keratitis, herpetic keratitis, conical cornea, dystrophia epithelialis corneae, corneal leukoma, ocular pemphigus, Mooren's ulcer, Scleritis, Graves' ophthalmopathy, Vogt-Koyanagi-Harada syndrome, sarcoidosis, etc.; pollen allergies, reversible obstructive airway disease, which includes condition such as asthma (for example, bronchial asthma, allergic asthma, intrinsic asthma, extrinsic asthma and dust asthma), particularly chronic or inveterate asthma (for example, late asthma and airway hyper-responsiveness), bronchitis and the like; inflammation of mucous and blood vessels such as gastric ulcers, vascular damage caused by ischemic diseases and thrombosis, ischemic bowel diseases, inflammatory bowel diseases, necrotizing enterocolitis, intestinal lesions associated with thermal burns and leukotriene B₄-mediated diseases; intestinal inflammations/allergies such as Coeliac diseases, proctitis, eosinophilic gastroenteritis, mastocytosis, Crohn's disease and ulcerative colitis; food-related allergic diseases which have symptomatic manifestation remote from the gastrointestinal tract (e.g. migraine,

- 36 -

rhinitis and eczema); renal diseases such as interstitial nephritis, Good-pasture's syndrome, hemolytic-uremic syndrome and diabetic nephropathy; nervous diseases such as multiple myositis, Guillain-Barre syndrome, Meniere's disease, polyneuritis, multiple neuritis,
5 mononeuritis and radiculopathy; endocrine diseases such as hyperthyroidism and Basedow's disease; hematic diseases such as pure red cell aplasia, aplastic anemia, hypoplastic anemia, idiopathic thrombocytopenic purpura, autoimmune hemolytic anemia, agranulocytosis, pernicious anemia, megaloblastic anemia and
10 anerythroplasia; bone diseases such as osteoporosis; respiratory diseases such as sarcoidosis, fibroid lung and idiopathic interstitial pneumonia; skin disease such as dermatomyositis, leukoderma vulgaris, ichthyosis vulgaris, photoallergic sensitivity and cutaneous T cell lymphoma; circulatory diseases such as arteriosclerosis, atherosclerosis, aortitis
15 syndrome, polyarteritis nodosa and myocardosis; collagen diseases such as scleroderma, Wegener's granuloma and Sjogren's syndrome; adiposis; eosinophilic fascitis; periodontal disease such as lesions of gingiva, periodontium, alveolar bone and substantia ossea dentis; nephrotic syndrome such as glomerulonephritis; male pattern alopecia
20 or alopecia senilis by preventing epilation or providing hair germination and/or promoting hair generation and hair growth; muscular dystrophy; Pyoderma and Sezary's syndrome; Addison's disease; active oxygen-mediated diseases, as for example organ injury such as ischemia-reperfusion injury of organs (such as heart, liver, kidney and digestive tract) which occurs upon preservation, transplantation or ischemic
25 disease (for example, thrombosis and cardiac infraction); intestinal diseases such as endotoxin-shock, pseudomembranous colitis and colitis caused by drug or radiation; renal diseases such as ischemic acute renal insufficiency and chronic renal insufficiency; pulmonary diseases such
30 as toxinoses caused by lung-oxygen or drug (for example, paracort and bleomycins), lung cancer and pulmonary emphysema; ocular diseases such as cataract, siderosis, retinitis, pigmentosa, senile macular degeneration, vitreal scarring and corneal alkali burn; dermatitis such as erythema multiforme, linear IgA bullous dermatitis and cement

- 37 -

dermatitis; and others such as gingivitis, periodontitis, sepsis, pancreatitis, diseases caused by environmental pollution (for example, air pollution), aging, carcinogenesis, metastasis of carcinoma and hypobaropathy; disease caused by histamine or leukotriene-C₄ release;

5 Behcet's disease such as intestinal-, vasculo- or neuro-Behcet's disease, and also Behcet's which affects the oral cavity, skin, eye, vulva, articulation, epididymis, lung, kidney and so on. Furthermore, the compounds of the invention are useful for the treatment and prevention of hepatic disease such as immunogenic diseases (for example, chronic

10 autoimmune liver diseases such as the group consisting of autoimmune hepatitis, primary biliary cirrhosis and sclerosing cholangitis), partial liver resection, acute liver necrosis (e.g. necrosis caused by toxin, viral hepatitis, shock, or anoxia), B-virus hepatitis, non-A/non-B hepatitis, cirrhosis (such as alcoholic cirrhosis) and hepatic failure such as

15 fulminant hepatic failure, late-onset hepatic failure and "acute-on-chronic" liver failure (acute liver failure on chronic liver diseases), and moreover are useful for various diseases because of their useful activity such as augmentation of chemotherapeutic effect, preventing or treating activity of cytomegalovirus infection, particularly HCMV infection, and

20 antiinflammatory activity; and

The compounds of the present invention may also be used in the treatment of immunodepression or a disorder involving immunodepression, such as AIDS, cancer, senile dementia, trauma (including wound healing, surgery and shock) chronic bacterial

25 infection, and certain central nervous system disorders.

A method of treating a condition in a mammal, the treatment of which is effected or facilitated by K_v1.3 inhibition, comprising the administration, in an amount that is effective at inhibiting K_v1.3, of a compound of Formula I. The method of treating

30 a condition in a mammal, the treatment of which is effected or facilitated by K_v1.3 inhibition, wherein the condition is selected from the group consisting of: immunemediated diseases such as the resistance by transplantation of organs or tissue such as heart, kidney, liver, medulla ossium, skin, cornea, lung, pancreas, intestinum tenue, limb,

- 38 -

muscle, nervus, duodenum, small-bowel, pancreatic-islet-cell, including xeno transplants, etc.; graft-versus-host diseases brought about by medulla ossium transplantation; autoimmune diseases such as rheumatoid arthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, type I diabetes uveitis, juvenile-onset or recent-onset diabetes mellitus, posterior uveitis, allergic encephalomyelitis, glomerulonephritis, and the like; and further infectious diseases caused by pathogenic microorganisms. Further uses may include the treatment and prophylaxis of inflammatory and hyperproliferative skin diseases and cutaneous manifestations of immunologically mediated illnesses, such as psoriasis, atopic dermatitis, contact dermatitis and further eczematous dermatitises and further eczematous dermatitises, seborrhoeis dermatitis, Lichen planus, Pemphigus, bullous pemphigoid, Epidermolysis bullosa, urticaria, angioedemas, vasculitides, erythemas, cutaneous eosinophilias, Lupus erythematosus, acne and Alopecia areata; various eye diseases (autoimmune and otherwise) such as keratoconjunctivitis, vernal conjunctivitis, uveitis associated with Behcet's disease, keratitis, herpetic keratitis, conical cornea, dystrophia epithelialis corneae, corneal leukoma, ocular pemphigus, Mooren's ulcer, Scleritis, Graves' ophthalmopathy, Vogt-Koyanagi-Harada syndrome, sarcoidosis, etc.; reversible obstructive airway disease, which includes condition such as asthma (for example, bronchial asthma, allergic asthma, intrinsic asthma, extrinsic asthma and dust asthma), particularly chronic or inveterate asthma (for example, late asthma and airway hyper-responsiveness), bronchitis and the like; inflammation of mucous and blood vessels such as gastric ulcers, vascular damage caused by ischemic diseases and thrombosis, ischemic bowel diseases, inflammatory bowel diseases, necrotizing enterocolitis, intestinal lesions associated with thermal burns and leukotriene B₄-mediated diseases; intestinal inflammations/allergies such as Coeliac diseases, proctitis, eosinophilic gastroenteritis, mastocytosis, Crohn's disease and ulcerative colitis; food-related allergic diseases which have symptomatic manifestation remote from the gastrointestinal tract (e.g. migraine, rhinitis and

- 39 -

- eczema); renal diseases such as interstitial nephritis, Good-pasture's syndrome, hemolytic-uremic syndrome and diabetic nephropathy; nervous diseases such as multiple myositis, Guillain-Barre syndrome, Meniere's disease, polyneuritis, multiple neuritis, mononeuritis and
- 5 radiculopathy; endocrine diseases such as hyperthyroidism and Basedow's disease; hematic diseases such as pure red cell aplasia, aplastic anemia, hypoplastic anemia, idiopathic thrombocytopenic purpura, autoimmune hemolytic anemia, agranulocytosis, pernicious anemia, megaloblastic anemia and anerythroplasia; bone diseases such as
- 10 osteoporosis; respiratory diseases such as sarcoidosis, fibroid lung and idiopathic interstitial pneumonia; skin disease such as dermatomyositis, leukoderma vulgaris, ichthyosis vulgaris, photoallergic sensitivity and cutaneous T cell lymphoma; circulatory diseases such as arteriosclerosis, atherosclerosis, aortitis syndrome, polyarteritis nodosa and
- 15 myocardosis; collagen diseases such as scleroderma, Wegener's granuloma and Sjogren's syndrome; adiposis; eosinophilic fascitis; periodontal disease such as lesions of gingiva, periodontium, alveolar bone and substantia ossea dentis; nephrotic syndrome such as glomerulonephritis; male pattern alopecia or alopecia senilis by
- 20 preventing epilation or providing hair germination and/or promoting hair generation and hair growth; muscular dystrophy; Pyoderma and Sezary's syndrome; Addison's disease; active oxygen-mediated diseases, as for example organ injury such as ischemia-reperfusion injury of organs (such as heart, liver, kidney and digestive tract) which occurs
- 25 upon preservation, transplantation or ischemic disease (for example, thrombosis and cardiac infraction): intestinal diseases such as endotoxin-shock, pseudomembranous colitis and colitis caused by drug or radiation; renal diseases such as ischemic acute renal insufficiency and chronic renal insufficiency; pulmonary diseases such as toxinoses caused
- 30 by lung-oxygen or drug (for example, paracort and bleomycins), lung cancer and pulmonary emphysema; ocular diseases such as cataract, siderosis, retinitis, pigmentosa, senile macular degeneration, vitreal scarring and corneal alkali burn; dermatitis such as erythema multiforme, linear IgA bullous dermatitis and cement dermatitis; and

- 40 -

others such as gingivitis, periodontitis, sepsis, pancreatitis, diseases caused by environmental pollution (for example, air pollution), aging, carcinogenesis, metastasis of carcinoma and hypobaropathy; disease caused by histamine or leukotriene-C₄ release; Behcet's disease such as

5 intestinal-, vasculo- or neuro-Behcet's disease, and also Behcet's which affects the oral cavity, skin, eye, vulva, articulation, epididymis, lung, kidney and so on. Furthermore, the compounds of the invention are useful for the treatment and prevention of hepatic disease such as immunogenic diseases (for example, chronic autoimmune liver diseases

10 such as the group consisting of autoimmune hepatitis, primary biliary cirrhosis and sclerosing cholangitis), partial liver resection, acute liver necrosis (e.g. necrosis caused by toxin, viral hepatitis, shock, or anoxia), B-virus hepatitis, non-A/non-B hepatitis, cirrhosis (such as alcoholic cirrhosis) and hepatic failure such as fulminant hepatic failure, late-

15 onset hepatic failure and "acute-on-chronic" liver failure (acute liver failure on chronic liver diseases), and moreover are useful for various diseases because of their useful activity such as augmentation of chemotherapeutic effect, preventing or treating activity of cytomegalovirus infection, particularly HCMV infection, and

20 antiinflammatory activity; and immunodepression or a disorder involving immunodepression, such as AIDS, cancer, senile dementia, trauma (including wound healing, surgery and shock), chronic bacterial infection, and certain central nervous system disorders.

An embodiment of the invention is a method for the

25 treatment of autoimmune diseases. Another embodiment of the invention is a method for the prevention of rejection of foreign organ transplants comprising administering to a patient in need of such treatment a therapeutically effective amount of a compound of formula I.

One end result of an autoimmune or a rejection process is

30 tissue destruction caused by inflammatory cells and the mediators they release. Anti-inflammatory agents such as NSAID's and corticosteroids act principally by blocking the effect or secretion of these mediators,

- 41 -

but do nothing to modify the immunologic basis of the disease. On the other hand, cytotoxic agents, such as cyclophosphamide, act in such a nonspecific fashion that both the normal and autoimmune responses are shut off. Indeed, patients treated with such nonspecific immuno-
5 suppressive agents are as likely to succumb from infection as they are from their autoimmune disease.

Cyclosporin A, which was approved by the US FDA in 1983, is currently the leading drug used to prevent rejection of transplanted organs. The drug acts by inhibiting the body's immune
10 system from mobilizing its vast arsenal of natural protecting agents to reject the transplant's foreign protein. Though cyclosporin A is effective in fighting transplant rejection, it is nephrotoxic and is known to cause several undesirable side effects including kidney failure, abnormal liver function and gastrointestinal discomfort.

15 Newer, safer drugs exhibiting fewer side effects are constantly being searched for in the field. The present invention provides for immunosuppressant agents which are inhibitors of a voltage dependent potassium channel, $K_v1.3$, that is found on human T-lymphocytes.

20 Potassium channels modulate a number of cellular events such as muscle contraction, neuro-endocrine secretion, frequency and duration of action potentials, electrolyte homeostasis, and resting membrane potential. These channels comprise a family of proteins that have been classified according to their biophysical and pharmacological
25 characteristics. Inhibition of K^+ channels, in their role as modulators of the plasma membrane potential in human T-lymphocytes, has been postulated to play a role in eliciting immunosuppressive responses. In regulating membrane potential, K^+ channels play a role in the regulation of intracellular Ca^{++} homeostasis, which has been found to
30 be important in T-cell activation. The biochemical characterization of K^+ channels is underdeveloped, due to the paucity of selective high affinity probes.

Functional voltage-gated K^+ channels can exist as multimeric structures formed by the association of either identical or

- 42 -

dissimilar subunits. This phenomena is thought to account for the wide diversity of K^+ channels. However, subunit compositions of native K^+ channels and the physiologic role that particular channels play are, in most cases, still unclear.

5 The $K_v1.3$ channel is a voltage-gated potassium channel that is found in neurons, blood cells, osteoclasts and T-lymphocytes. The Chandy and Cahalan laboratories proposed a hypothesis that blocking the $K_v1.3$ channel would elicit an immunosuppressant response. (Chandy *et al.*, *J. Exp. Med.* 160, 369, 1984; Decoursey *et al.*,
10 Nature, 307, 465, 1984). However, the K^+ channel blockers employed in their studies were non-selective. Until research with the peptide margatoxin, a peptide found in scorpion venom, no specific inhibitor of the $K_v1.3$ channel existed to test this hypothesis. Although a laboratory (Price *et al.*, *Proc. Natl. Acad. Sci. USA*, 86, 10171, 1989) showed that
15 charybdotoxin would block $K_v1.3$ in human T cells, charybdotoxin was subsequently shown to inhibit four different K^+ channels ($K_v1.3$ and three distinct small conductance Ca^{++} activated K^+ channels) in human T-lymphocytes, limiting the use of this toxin as a probe for the physiological role of $K_v1.3$ (Leonard *et al.*, *Proc. Natl. Acad. Sci. USA*,
20 89, 10094, 1992). Margatoxin, on the other hand, blocks only $K_v1.3$ in T-cells, and has immunosuppressant activity in both in vitro and in vivo models. (Lin *et al.*, *J. Exp. Med.* 177, 637, 1993). Since the compounds of the embodiments of this invention produce blockade of $K_v1.3$, they will also inhibit T-cell activation.

25 Also within the scope of this invention is a method of treating a condition in a mammal, the treatment of which is effected or facilitated by $K_v1.3$ inhibition, comprising the administration of a pharmaceutical composition comprising a suitable pharmaceutical carrier and a compound of Formula (I), in an amount that is effective at
30 inhibiting $K_v1.3$.

 Also within the scope of this invention is a combination therapy comprising a compound of formula I and one or more immunosuppressant agents. These immunosuppressant agents within the scope of this invention include, but are not limited to, IMUREK®

- 43 -

azathioprine sodium, brequinar sodium, SPANIDIN® gusperimus trihydrochloride (also known as deoxyspergualin), mizoribine (also known as bredinin), CELLCEPT® mycophenolate mofetil, NEORAL® Cyclosporin A (also marketed as different formulation of Cyclosporin A under the trademark SANDIMMUNE®), PROGRAF® tacrolimus (also known as FK-506) and RAPIMMUNE® sirolimus (also known as rapamycin), leflunomide (also known as HWA-486), glucocorticoids, such as prednisolone and its derivatives, antibody therapies such as orthoclone (OKT3) and Zenapax and antithymocyte globulins, such as thymoglobulins.

Using the methodologies described below, representative compounds of the invention were evaluated and found to exhibit IC50 values of at least $<10 \mu\text{M}$ in any of the assays thereby demonstrating and confirming the utility of the compounds of the invention as Kv1.3 inhibitors and immunosuppressants.

T CELL IL-2 ASSAY

Peripheral blood mononuclear (MNC) cells from healthy donors were separated by density centrifugation with ficoll-hypaque (LSM, Organon Teknika, Durham, NC), followed by rosetted with neuraminidase treated sheep red blood cells (SRBC). After another centrifugation with leucocyte separation medium (LSM), the SRBC of the rosetted T cells were then lysed with ammonium chloride lysing buffer (GIBCO, Grand Island, NY). Such purified T cells were resuspended at 3×10^6 /ml in RPMI 1640 culture medium (GIBCO) supplemented with 10% fetal calf serum (Sigma, St. Louis, MO), 100 mM glutamine, 1 mM sodium pyruvate, 0.1 mM non-essential amino acids, and 1 % penn-strep (GIBCO). The cell suspension was immediately distributed into 96 well round-bottom microculture plates (Costar) at 200 μl /well. The various dilutions of test compound were then added in triplicate wells at 25 μl /well, incubated for 30 min at 37°C. Ionomycin (125 ng/ml), and PMA (1 or 5 ng/ml), were added to the appropriate wells. The culture plates were then incubated at 37°C in a humidified atmosphere of 5% CO₂ - 95% air for 18-24 hours. The

- 44 -

supernatants were removed, and assayed for IL-2 with an IL-2 capture ELISA, using monoclonal anti-IL-2, and biotinylated goat anti-IL-2 antibodies (unconjugated antibodies purchased from R&D System, Minneapolis, MN). The ELISA was developed with streptavidin
5 conjugated peroxidase (Zymed, San Francisco, CA) and substrate for peroxidase (Sigma). Mean OD and units of IL-2 of the replicate wells were calculated from standard curve, created with recombinant IL-2 (Collaborative Biomedical Products, Bedford, MA) and the results were expressed as concentration of compound required to inhibit IL-2
10 production of T cells by 50%.

T CELL PROLIFERATION ASSAY

Peripheral blood mononuclear cells (MNC) from healthy
15 donors were separated by density centrifugation with ficoll-hypaque (LSM, Organon Teknika, Durham, NC). After washing the MNC with complete media (RPMI 1640 medium with 5% fetal calf serum, 100 mM glutamine, 1 mM sodium pyruvate, 0.1 mM non-essential amino acid, and 1% penn-strep, obtained from GIBCO, Grand Island, NY), they
20 were then irradiated at 7500 RADS, and resuspended at $4-4.5 \times 10^6$ cells/ml in complete media. Another aliquot of MNC were rosetted with neuraminidase treated SRBC. After another centrifugation with LSM, the sheep red blood cells (SRBC) of these rosetted T cells were then lysed with ammonium chloride lysing buffer (GIBCO, Grand
25 Island, NY). After washing 2X with complete media, these purified T cells were also resuspended at $2-2.5 \times 10^6$ cells/ml in complete media. The various dilutions of the compound were added in triplicates at 50 ul/well of a 96 well flat-bottom microculture plate (Costar, Cambridge, MA). T cell suspension was then immediately distributed into the wells
30 at 100 ul/well. After incubating the cells with compound for 30 min. at 37°C in a humidified atmosphere of 5% CO₂ - 95% air, 20 µl/well of anti-CD3 (Ortho Diagnostic, NJ) at final conc. of 0.3 ng/ml was added, followed by 50 µl of the irradiated MNC. The culture plates were then incubated at 37°C in a humidified atmosphere of 5% CO₂ - 95% air for
35 72 hours. The proliferation of T lymphocytes was assessed by

- 45 -

measurement of tritiated thymidine incorporation. During the last 18-24 hrs. of culturing, the cells were pulse-labeled with 2 μ Ci/well of tritiated thymidine (NEN, Cambridge, MA). The cultures were harvested on glass fiber filters using a multiple sample harvester (MACH-II, Wallac, Gaithersburg, MD). Radioactivity of filter discs corresponding to individual wells was measured by standard liquid scintillation counting methods (Betaplate Scint Counter, Wallac). Mean counts per minute of replicate wells were calculated and the results were expressed as concentration of compound required to inhibit tritiated thymidine uptake of T cells by 50%.

KV1.3-RUBIDIUM EFFLUX ASSAY

CHO cells transfected with Kv1.3 channels at site densities of approximately 40,000 sites/cell are plated into 96 well culture plates and maintained in Iscove's Modified Dulbecco's Medium (IMDM, with L-Glutamine and HEPES, JRH Biosciences). Cells are incubated overnight with $^{86}\text{Rb}^+$ (3 μ Ci/ml, Dupont-NEN) in the glutamine supplemented IMDM. After aspiration of the media, 100 μ l of Low K Buffer (in mM, 6.5 KCl, 125 NaCl, 1 CaCl_2 , 2 MgCl_2 , 10 HEPES, pH adjusted to 7.2 with NaOH) is added to each well followed by 100 μ l test samples in Low K Buffer also containing 0.2% BSA and 2 mM ouabain. Samples are tested at either 1 μ g/ml for routine screening or at a variety of concentrations encompassing at least 1/10 to 10 times the putative IC₅₀ of test compound to determine potency. After a fixed preincubation time, which is usually 10 min, the samples are aspirated. The Kv1.3 channels are opened by depolarization of the cells with High K Buffer (final concentrations, in mM, 63.25 KCl, 68.25 NaCl, 1 CaCl_2 , 2 MgCl_2 , 10 HEPES, pH adjusted to 7.2 with NaOH) also containing test compounds. To measure $^{86}\text{Rb}^+$ efflux through the channels, aliquots of 100 μ l are taken from each well after a given time and added to plates containing 100 μ l MicroScint-40 (Packard) for counting by liquid scintillation techniques. MicroScint-40 (100 μ l) is then added to each well of the cell plate to determine the remaining $^{86}\text{Rb}^+$ activity. The efflux counts are normalized for the total amount of

- 46 -

$^{86}\text{Rb}^+$ that was in the cells by adding the efflux counts to the cell plate counts. Activity is determined by % inhibition of the efflux window that is established using a saturating concentration of margatoxin (MgTX), a 39 amino acid peptide that is a potent blocker of $\text{K}_v1.3$ channels ($\text{IC}_{50} =$
5 100 pM).

DOSAGE FORMS

As an immunosuppressive, these compounds are useful in
10 the treatment of autoimmune diseases, the prevention of rejection of foreign organ transplants and/or related afflictions, diseases and illnesses.

The compounds of this invention can be administered for the treatment of autoimmune diseases, the prevention of rejection of
15 foreign organ transplants and/or related afflictions, diseases and illnesses according to the invention by any means that effects contact of the active ingredient compound with the site of action in the body of a warm-blooded animal. For example, administration, can be oral, topical, including transdermal, ocular, buccal, intranasal, inhalation,
20 intravaginal, rectal, intracisternal and parenteral. The term "parenteral" as used herein refers to modes of administration which include subcutaneous, intravenous, intramuscular, intraarticular injection or infusion, intrasternal and intraperitoneal.

The compounds can be administered by any conventional
25 means available for use in conjunction with pharmaceuticals, either as individual therapeutic agents or in a combination of therapeutic agents. They can be administered alone, but are generally administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

30 For the purpose of this disclosure, a warm-blooded animal is a member of the animal kingdom possessed of a homeostatic mechanism and includes mammals and birds.

The dosage administered will be dependent on the age, health and weight of the recipient, the extent of disease, kind of
35 concurrent treatment, if any, frequency of treatment and the nature of

- 47 -

the effect desired. Usually, a daily dosage of active ingredient compound will be from about 1-500 milligrams per day. Ordinarily, from 10 to 100 milligrams per day in one or more applications is effective to obtain desired results. These dosages are the effective amounts for the treatment of autoimmune diseases, the prevention of rejection of foreign organ transplants and/or related afflictions, diseases and illnesses.

The active ingredient can be administered orally in solid dosage forms, such as capsules, tablets, troches, dragées, granules and powders, or in liquid dosage forms, such as elixirs, syrups, emulsions, dispersions, and suspensions. The active ingredient can also be administered parenterally, in sterile liquid dosage forms, such as dispersions, suspensions or solutions. Other dosages forms that can also be used to administer the active ingredient as an ointment, cream, drops, transdermal patch or powder for topical administration, as an ophthalmic solution or suspension formation, i.e., eye drops, for ocular administration, as an aerosol spray or powder composition for inhalation or intranasal administration, or as a cream, ointment, spray or suppository for rectal or vaginal administration.

Gelatin capsules contain the active ingredient and powdered carriers, such as lactose, starch, cellulose derivatives, magnesium stearate, stearic acid, and the like. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric coated for selective disintegration in the gastrointestinal tract.

Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance.

In general, water, a suitable oil, saline, aqueous dextrose (glucose), and related sugar solutions and glycols such as propylene glycol or polyethylene glycols are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably contain a

- 48 -

water soluble salt of the active ingredient, suitable stabilizing agents, and if necessary, buffer substances. Antioxidizing agents such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, are suitable stabilizing agents. Also used are citric acid and its salts and sodium EDTA. In addition, parenteral solutions can contain
5 preservatives, such as benzalkonium chloride, methyl- or propylparaben, and chlorobutanol.

Suitable pharmaceutical carriers are described in
10 *Remington's Pharmaceutical Sciences*, A. Osol, a standard reference text in this field.

For administration by inhalation, the compounds of the present invention may be conveniently delivered in the form of an aerosol spray presentation from pressurized packs or nebulisers. The compounds may also be delivered as powders which may be formulated
15 and the powder composition may be inhaled with the aid of an insufflation powder inhaler device. The preferred delivery system for inhalation is a metered dose inhalation (MDI) aerosol, which may be formulated as a suspension or solution of a compound of Formula I in suitable propellants, such as fluorocarbons or hydrocarbons.

20 For ocular administration, an ophthalmic preparation may be formulated with an appropriate weight percent solution or suspension of the compounds of Formula I in an appropriate ophthalmic vehicle, such that the compound is maintained in contact with the ocular surface for a sufficient time period to allow the compound to penetrate the
25 corneal and internal regions of the eye.

Useful pharmaceutical dosage-forms for administration of the compounds of this invention can be illustrated as follows:

CAPSULES

30

A large number of unit capsules are prepared by filling standard two-piece hard gelatin capsules each with 100 milligrams of powdered active ingredient, 150 milligrams of lactose, 50 milligrams of cellulose, and 6 milligrams magnesium stearate.
35

- 49 -

SOFT GELATIN CAPSULES

5 A mixture of active ingredient in a digestible oil such as soybean oil, cottonseed oil or olive oil is prepared and injected by means of a positive displacement pump into gelatin to form soft gelatin capsules containing 100 milligrams of the active ingredient. The capsules are washed and dried.

TABLETS

10

A large number of tablets are prepared by conventional procedures so that the dosage unit is 100 milligrams of active ingredient, 0.2 milligrams of colloidal silicon dioxide, 5 milligrams of magnesium stearate, 275 milligrams of microcrystalline cellulose, 11
15 milligrams of starch and 98.8 milligrams of lactose. Appropriate coatings may be applied to increase palatability or delay absorption.

INJECTABLE

20

A parenteral composition suitable for administration by injection is prepared by stirring 1.5% by weight of active ingredient in 10% by volume propylene glycol. The solution is made to volume with water for injection and sterilized.

25

SUSPENSION

An aqueous suspension is prepared for oral administration so that each 5 milliliters contain 100 milligrams of finely divided active ingredient, 100 milligrams of sodium carboxymethyl cellulose, 5
30 milligrams of sodium benzoate, 1.0 grams of sorbitol solution, U.S.P., and 0.025 milliliters of vanillin.

The same dosage forms can generally be used when the compounds of this invention are administered stepwise or in conjunction with another therapeutic agent. When drugs are administered in
35 physical combination, the dosage form and administration route should be selected depending on the compatibility of the combined drugs. Thus


- 50 -

the term coadministration is understood to include the administration of the two agents concomitantly or sequentially, or alternatively as a fixed dose combination of the two active components.

- The following examples illustrate the preparation of the compounds of Formula I and as such are not to be considered as limiting the invention set forth in the claims appended hereto.

EXAMPLE 1

- 10 A Method Of Extracting The Compounds Of Formula 1(a) and 1(b)
From *Spachea correa*



Formula 1(a) b is a single bond and R is OAc

Formula 1(b) b is a double bond and R is OAc

- One gram of an ethanol extract of the roots of *Spachea correa* was partitioned between 100 ml of hexane (twice) and 100 ml of 90% aqueous methanol. After separation of the phases, the defatted methanol was concentrated down under vacuum to give an aqueous suspension. This was diluted out to 100 ml with water and extracted, with 100 ml of methylene chloride.

- The bioactive methylene chloride extract was dried down to give 12 mg of residue. This was first fractionated by preparative thin layer chromatography (TLC) on a 20 cm by 20 cm E. Merck silica gel 60F254 plate of 1mm thickness using methylene chloride-ethyl acetate 1:1 (v/v) as solvent, then by high performance liquid chromatography (HPLC) using a Zorbax RxC8 4.6 mm x 25 cm column, operated at 50°C and eluted with a 50 minute gradient of acetonitrile:water (1:1,

- 51 -

v/v) to 100% acetonitrile, delivered at 1 ml/min, to afford 4 mg of compound 1(a) and 1 mg of 1(b).

Homogeneity of the preparations was ascertained in several TLC systems, such as E. Merck silica gel 60F254, methylene chloride-ethyl acetate 1:1, Rf 1(a) 0.4, Rf 1(b) 0.3; Whatman KC18, methanol-water 9:1, Rf 1(a) 0.65, Rf 1(b) 0.75 and by HPLC using a Zorbax Rx C8 column, acetonitrile-water 3:2, k' 1(a) 4.15, k' 1(b) 3.30; and by NMR.

Mass spectra were recorded on JEOL SX-102A (electron impact, EI, 903V) and JEOL HX110 (Fast Atom Bombardment, FAB) mass spectrometers. Exact mass measurements were performed at high resolution (HR-EI) using perfluorokerosene (PFK) as the internal standard. Trimethylsilyl derivatives were prepared with a 1:1 mixture of BSTFA-pyridine at room temperature. The FAB spectrum was run in a matrix of dithiothreitol dithioerthritol (20/80).

The compound of Formula 1(a) runs underivatized by EI. The molecular ion is observed a m/z 788 and three successive losses of acetic acid are observed. The base peak is observed a m/z 334. The compound does not silylate. Scanning HR-EI indicated a molecular formula of C₄₀H₅₂O₁₆. A table of the critical HR-EI data is given below.

	<u>Observed m/z</u>	<u>Formula</u>	<u>Assignment</u>
	788.3220	C ₄₀ H ₅₂ O ₁₆	M ⁺
25	728.3040	C ₃₈ H ₄₈ O ₁₄	M-acetic acid
	668.2834	C ₃₆ H ₄₄ O ₁₂	M-2 x acetic acid
	334.1417	C ₁₈ H ₂₂ O ₆	base peak

¹³C NMR spectra were recorded for the compound of Formula 1(a) in CD₂Cl₂ at 100 MHz on a Varian Unity 400 NMR spectrometer at 20°C. Chemical shifts are given in ppm relative to tetramethylsilane (TMS) at zero ppm using the solvent peak at 53.8 ppm as internal standard. The following data were observed: 15.0, 15.2, 16.8, 17.1, 20.7*, 20.9, 21.1, 21.6, 21.8, 22.2, 35.6, 40.8*, 42.1, 43.6,

- 52 -

45.1, 47.5, 49.3*, 53.5, 59.1, 62.6, 63.5, 66.1, 66.7*, 68.4*, 69.9, 73.9, 75.0, 75.6, 77.1*, 119.4, 123.7, 138.9, 143.0, 167.7, 169.2, 169.3*, 170.25, 170.31, 170.8, 171.3 ppm (where the * signifies the observation as broad resonances). The carbon count of 40 is in agreement with the
 5 molecular formula $C_{40}H_{52}O_{16}$ derived by scanning HR EI-MS.

The 1H NMR spectra of compound of Formula(a) is provided as Figure 1. The spectra was recorded at 400 MHz in CD_2Cl_2 on a Varian Unity 400 NMR spectrometer at 25°C. Chemical shifts are in ppm relative to TMS at zero ppm using the solvent peak at δ 5.32 as
 10 the internal standard.


The mass spectra of the compound of Formula 1(b) was obtained as above. The following results were obtained.

	<u>Observed m/z</u>	<u>Formula</u>	<u>Assignment</u>
15	786.3075	$C_{40}H_{50}O_{16}$	M^+
	726.2886	$C_{38}H_{46}O_{14}$	M-acetic acid
	666.2651	$C_{36}H_{42}O_{12}$	M-2 x acetic acid
	606.2451	$C_{34}H_{38}O_{10}$	M-3 x acetic acid
	489.2099	$C_{26}H_{33}O_9$	base peak
20	471.1992	$C_{26}H_{31}O_8$	

^{13}C NMR spectra were recorded for the compound of Formula 1(b) using the procedure described above. The following results were observed: 14.8, 14.9, 17.3, 20.8, 20.9, 21.3, 21.7, 21.8,
 25 21.9, 27.1, 35.1, 40.6, 42.3, 45.4, 48.1, 50.4, 53.5, 54.1, 57.8, 63.7, 66.2, 67.8, 68.6, 71.4, 73.3, 73.8, 74.4, 119.5, 121.1, 124.3, 137.1, 138.9, 143.3, 167.6, 168.6, 169.3, 169.5, 169.9, 171.0, 171.7 ppm.

The carbon count of 40 is in agreement with the molecular formula $C_{40}H_{50}O_{16}$ derived by scanning HR EI-MS.

- 53 -

EXAMPLE 2A Method Of Extracting The Compounds Of Formula 1(c) And 1(d)
From *Spachea Correa*

Formula 1(c) b is a single bond and R is OH

Formula 1(d) b is a double bond and R is OH

5

10 Analogs of the compounds of Formula 1(a) and 1(b) could be detected in the crude extract and fractions thereof when the process of Example 1 was carried out on a larger scale. Thus, 50 g of ethanol extract were partitioned as described in Example 1 using 900 ml of each solvent at each step.

15 Partial purification of the methylene chloride extract was achieved by column chromatography on E. Merck silica gel 60 (120 ml), eluting with a step gradient of ethyl acetate in methylene chloride. The step gradient was designed so that the column was washed first with 100% methylene chloride and then with methylene chloride-ethyl acetate mixtures of 9:1, 8:2, 3:2, 2:1, 1:1, 1:2, 2:8 and 1:9. Ultimately the column was washed with 100% ethyl acetate. Fractions eluted with methylene chloride-ethyl acetate 3:2 were enriched in compound of Formula 1(a) and 1(b). These were resolved by HPLC using a Zorbax
20 RxC8 9 mm x 25 cm column, maintained at 50°C and eluted at 4 ml/min with acetonitrile-water 1:1 v/v. Three identical runs finally afforded 100 mg and 20 mg respectively of 1(a) and 1(b) after crystallization from methanol. Later-eluting fractions from the silica gel column above were found to contain at least two related compounds based on
25 UV spectra and color reactions on TLC plates. Material from the

- 54 -

methylene chloride-ethyl acetate 1:1 and 1:2 washings were combined and evaporated down. Separation was achieved on the same HPLC column as above, eluting with a 50 minute gradient of 30% to 50% acetonitrile in water. Two identical runs gave 6 mg of purified compound 1(c). Fractions containing the compound of Formula 1(d) were again processed by HPLC (same column) using acetonitrile-water 3:7 delivered isocratically, to yield 2 mg of purified Formula 1(d).

The mass spectra of these compounds were recorded on a Finnigan TSQ700 mass spectrometer (electrospray ionization, ESI). The samples were analyzed by LC/MS using a 2.1x150mm C8 column at 0.2ml/min. with a mobile phase of 45% acetonitrile/0.01M aqueous ammonium acetate at 50°C. Component 1(d) had a retention time of 10.5 min. and a molecular weight of 744 which is observed a m/z: 745 (M⁺H), 762 (M⁺NH₃), 786 (M + H + MeCN). Component 1(c) has a retention time of 11.8 and a molecular weight of 746 which is observed at m/z: 747 (M⁺H), 764 (M⁺NH₃) and 788 (M + H + MeCN).

The ¹³C NMR spectra obtained for the compound of Formula 1(c) using the conditions previously described is as follows: 15.1 (2x), 16.9, 19.8, 20.8, 20.91, 20.94, 21.9, 22.3, 35.6, 40.6, 42.2, 43.9, 45.0, 47.7, 50.8, 53.5, 55.6, 61.8, 63.5, 66.0, 67.6 (2x), 69.8, 70.0, 73.9, 75.0, 75.6, 119.3, 123.7, 139.0, 144.4, 167.8, 169.2, 169.5, 170.1, 170.4, 171.4 ppm.

The carbon count of 38 is in agreement with the molecular formula C₃₈H₅₀O₁₆ derived by scanning HR EI-MS.

EXAMPLE 3

Separation By HPLC

Compounds of this invention were characterized by the following behavior during HPLC separation on a Zorbax RxC8 4.6 mm x 25 cm column, maintained at 50°C and eluted at 1 ml/min with acetonitrile-water 3:2 v/v):

Compound 1(a): k' = 4.15; 1(b): k'=3.30; 1(c): k'=2.30; 1(d): k'=2.10.

- 55 -

Analyses using this HPLC system can be used to quantify the compounds in the crude extract or other mixtures, by comparing the absorbance of HPLC peaks at a wavelength of 220 nm with that produced by injections of known (weighed) amounts of pure standards.

5

EXAMPLE 4

Additional Purification Procedure


10 A simplified purification process allows for rapid fractionation of even larger amounts of crude extract and the preparation of gram amounts of the compounds of Formula 1(a) and 1(b).

The ethanol extract is first dissolved at 20 grams per 150 ml in methanol. This solution is diluted with 150 ml of water and
15 then extracted three times with methylene chloride using 150 ml of methylene chloride each time. The pooled methylene chloride extracts are evaporated down and fractionation proceeds by repeated column chromatography on silica gel. One employs methylene chloride-methanol 97:3 in a first step; the mixed compounds of Formula 1(a) and
20 1(b) thus obtained are resolved by chromatographing on fresh silica gel eluted with methylene chloride-ethyl acetate 3:1. Volume of elution for the compound of Formula 1(a) ranges from about 2 to about 3.5 column volumes of solvent; that for the compound of Formula 1(b) is about 3 to about 4.5 column volumes. Finally, advantage is taken of the low
25 solubility of these compounds, and, after total resolution by chromatography, the compounds of Formula 1(a) and 1(b) can be precipitated and or crystallized from concentrated methanol solutions.

- 56 -

EXAMPLE 5

4,6,7,15,16-Pentakis(acetyloxy)-18-hydroxy-22-methoxycarbonyl-
 [6 α ,7 α ,15 β ,16 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-1,
 5 20(29),21-trien-3-one




A solution of 233 mg (0.56 mmole) of tungsten
 hexachloride in 8 mL of dry tetrahydrofuran was cooled to -78°C under
 10 nitrogen. Then 0.70 mL (1.12 mmole) of 1.6 M butyllithium was added
 and the solution was allowed to warm to room temperature over 30
 min. Then a solution of 111 mg (0.141 mmole) of 4,6,7,15,16-
 pentakis(acetyloxy)-21,22-epoxy-18-hydroxy-22-methoxycarbonyl-
 [6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-dinor-24-
 15 oxaoleana-1,20(29)-dien-3-one in 2 mL of dry tetrahydrofuran was
 added and the solution was heated under nitrogen at 55°C for 14 h. The
 mixture was applied to a 10 cm column of silica gel, which was washed
 with 2:1 ethyl acetate-hexane. The eluate was concentrated and purified
 by silica gel chromatography with 2:1 ethyl acetate-hexane to afford 95
 20 mg (88%) of the title compound as a white solid; ¹H NMR (CDCl₃) δ
 7.10 (s, 1H, C21-H); Mass Spectrum (APCI) 790 (M⁺NH₄).


- 57 -

EXAMPLE 6

4-(2-Bromobenzoyl)oxy-6,7,15,16-tetrakis(acetyloxy)-18-hydroxy-22-methoxycarbonyl[6 α ,7 α ,15 β ,16 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-1,20(29),21-trien-3-one



Step A: 6,7,15,16-Tetrakis(acetyloxy)-21,22-epoxy-4, 18-dihydroxy-22-methoxycarbonyl-[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-1,20(29)-dien-3-one



A solution of 102.1 mg (0.130 mmole) of 4,6,7,15,16-pentakis(acetyloxy)-21,22-epoxy-18-hydroxy-22-methoxycarbonyl-


- 58 -

[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-1,20(29)-dien-3-one in 4 mL of tetrahydrofuran and 2 mL of 3M aqueous HCl was heated at 40°C for 24h. The solution was diluted with dichloromethane and the layers were separated. The organic layer
 5 was washed with 0.1M phosphate buffer (pH 7), then was dried over MgSO₄ and concentrated. The residue was purified by silica gel chromatography with 2:1 ethyl acetate-hexane to afford 44.9 mg of the title compound as a white solid (46%); ¹H NMR (CDCl₃) δ 4.20 (q, 1H, J = 4.3 Hz, C4-H); Mass Spectrum (APCI): m/e 764 (M⁺NH₄).

10

Step B: 4-(2-Bromobenzoyl)oxy-6,7,15,16-tetrakis(acetyloxy)-21,22-epoxy-18-hydroxy-22-methoxycarbonyl-[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-1,20(29)-dien-3-one

15




To a solution of 17.5 mg (23.5 μ mole) of 6,7,15,16-tetrakis(acetyloxy)-21,22-epoxy-4,18-dihydroxy-22-methoxycarbonyl-[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-1,20(29)-dien-3-one in 0.5 mL pyridine was added 27.5 mL
 20 (237 μ mole) of benzoyl chloride. The solution was stirred at room temperature for 4 h, then was concentrated under reduced pressure.

- 59 -

The residue was first filtered through a plug of silica gel and then purified by HPLC (Waters RCM, μ Porosil, 10 mm X 10 cm) using a mixture of 9.6:6 (5:4:1 hexane-methyl *tert*-butyl ether-acetonitrile:hexane) to afford 17.3 mg (88%) of the title compound as a white solid; ^1H NMR δ 5.67 (1H, C4-H), 7.40-7.43 (m, 2H), 7.72 (dd, 1H, $J = 2.2, 6.9$ Hz), 7.78 (dd, 1H, $J = 2.3, 6.9$ Hz); Mass Spectrum (APCI): m/e 946. 948 ($^{79}\text{Br-M}^+\text{NH}_4$, $^{81}\text{Br-M}^+\text{NH}_4$).

10 Step C: 4-(2-Bromobenzoyl)oxy-6,7,15,16-tetrakis(acetyloxy)-18-hydroxy-22-methoxycarbonyl-[6 α ,7 α ,15 β ,16 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-1,20(29),21-trien-3-one



15 A solution of 218 mg (0.55 mmole) of tungsten hexachloride in 8 mL of dry tetrahydrofuran was cooled to -78°C under nitrogen. Then 0.69 mL (1.10 mmole) of 1.6M butyllithium was added and the solution was allowed to warm to room temperature over 30 min. Then a solution of 50.2 mg (0.141 mmole) of 4-(2-bromobenzoyloxy)-6,7,15,16-tetrakis(acetyloxy)-21,22-epoxy-18-hydroxy-22-methoxycarbonyl-[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-1,20(29)-dien-3-one in 2 mL of dry tetrahydrofuran was added and the solution was heated under nitrogen at

20


- 60 -

55°C for 14 h. The mixture was washed with 2.5 MNaOH and brine, then was dried over MgSO₄. The filtrate was purified by HPLC (Waters RCM, μ Porosil, 25mm x 10cm) using a 8.9:4 mixture of (5:4:1 hexane:methyl tert-butyl ether:acetonitrile):hexane to afford 31.3 mg (64%) of the title compound as a white solid; ¹H NMR (CDCl₃) δ 5.67 (1H, C4-H), 7.10 (s, 1H, C21-H), 7.40-7.43 (m, 2H), 7.72 (dd, 1H, J = 2.2, 6.9 Hz), 7.78 (dd, 1H, J = 2.3, 6.9 Hz); Mass Spectrum (APCI) m/e 930, 932 (⁷⁹Br-M+NH₄, ⁸¹Br-M+NH₄).

10

EXAMPLE 7

4,6,7,15,16-Pentakis(acetyloxy)-21,22-epoxy-18-hydroxy-22-methoxycarbonyl[6 α ,7 α ,15 β ,16 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-1,20(29),21-triene




15

Step A: 4,6,7,15,16-Pentakis(acetyloxy)-21,22-epoxy-18-hydroxy-22-methoxycarbonyl[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-1,20(29)-dien-3-ol

20

- 61 -



A solution of 3.0 g (3.8 mmole) of 4,6,7,15,16-pentakis(acetyloxy)-21,22-epoxy-18-hydroxy-22-methoxycarbonyl-[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-1,20(29)-dien-3-one in 20 mL of dry dichloromethane was cooled to 0°C under nitrogen. Then 9 mL of a 1M solution of lithium tri-(tert-butoxy)aluminum hydride was added dropwise and the solution was stirred at 0°C. After 18 h, the reaction was quenched by dropwise addition of 20 mL of 2M aqueous H₂SO₄ and the mixture was diluted


5
10
15

with 200 mL of ether. The layers were separated and the aqueous layer was washed with two 100 mL portions of ether. The organic layers were sequentially washed with 20 mL of 2M aqueous H₂SO₄ and brine, then were combined, dried over MgSO₄, and concentrated to afford 2.9 g (99%) of the title compound, which was used directly in the next step.

- 62 -

Step B: 4,6,7,15,16-Pentakis(acetyloxy)-21,22-epoxy-18-hydroxy-22-methoxycarbonyl[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-1,20(29)-diene

5




A sample of 2.9 g of crude 4,6,7,15,16-pentakis(acetyloxy)-21,22-epoxy-18-hydroxy-22-methoxycarbonyl-[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-1,20(29)-dien-3-ol was dissolved in 10 mL of dry dichloromethane under nitrogen. To this was added 10 mL of triethylsilane, and the solution was stirred at room temperature for 10 min. Then 2 mL (20 mmole) of boron trifluoride etherate was added and the mixture was stirred at room temperature for 15 min. The reaction was quenched by addition of 10 mL of saturated aqueous KHCO₃ solution and the resulting mixture was partitioned between ether and water. The water layer was washed with ether and the organic extracts were washed with brine, then were combined, dried over MgSO₄, and concentrated. The residue was purified by chromatography on silica gel using 30% ethyl acetate-hexane to afford 2.13 g (72%) of the title compound as a white solid; ¹H NMR (CDCl₃) δ 4.14, 4.34 (dd, AB, 2H, J = 12 Hz, C3-H); Mass Spectrum (APCI) m/e 792 (M⁺NH₄).

- 63 -

Step C: 4,6,7,15,16-Pentakis(acetyloxy)-18-hydroxy-22-methoxycarbonyl[6 α ,7 α ,15 β ,16 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-1,20(29),21-triene

5




A solution of 119 mg (0.300 mmole) of tungsten hexachloride in 4 mL of dry tetrahydrofuran was cooled to -78°C under nitrogen. Then 0.38 mL (0.61 mmole) of 1.6M butyllithium was added and the solution was allowed to warm to room temperature over 30 min. A solution of 34.6 mg (0.045 mmole) of 4,6,7,15,16-pentakis(acetyloxy)-21,22-epoxy-18-hydroxy-22-methoxycarbonyl-[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-1,20(29)-diene in 2 mL of dry tetrahydrofuran was added and the solution was heated under nitrogen at 50°C for 14 h. The mixture was diluted with 20 mL ether, and the mixture was washed with 2.5 M NaOH and brine, dried over MgSO₄, and concentrated. The residue was purified by HPLC (Waters RCM, μ Porosil, 25mm x 10cm) using a 6.75:7.0 mixture of (5:4:1 hexane:methyl tert-butyl ether:acetonitrile):hexane to afford 27 mg (80%) of the title compound as a white solid; ¹H NMR (CDCl₃) δ (s, 1H, C21-H); Mass Spectrum (APCI) 776 (M⁺NH₄).

- 64 -

EXAMPLE 8

4,6,7,15,16-Pentakis(acetyloxy)-21,22-epoxy-18-hydroxy-22-methoxycarbonyl[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-20(29)-en-3-one




As described in Scheme I, 4,5,6,15,16-pentakis(acetyloxy)-21,22-epoxy-18-hydroxy-22-methoxycarbonyl[6 α , 7 α , 15 β , 16 β , 21 β , 22 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-1,20(29)-diene-3-one, isolated from *Spachea correa* in liquid ammonia with lithium metal will result in the reduction of the C1 olefin group to produce the saturated lactone.

EXAMPLE 9

4,6,7,15,16-Pentakis(acetyloxy)-18-hydroxy-22-methoxycarbonyl-[6 α ,7 α ,15 β ,16 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-20(29),21-dien-3-one

- 65 -




A solution of 233 mg (0.56 mmole) of tungsten
 hexachloride in 8 mL of dry tetrahydrofuran is cooled to -78°C under
 5 nitrogen. Then 0.70 mL (1.12 mmole) of 1.6M butyllithium was added
 and the solution is allowed to warm to room temperature over 30 min.
 Then a solution of 111 mg (0.141 mmole) of 4,6,7,15,16-
 pentakis(acetyloxy)-21,22-epoxy-18-hydroxy-22-
 methoxycarbonyl[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-
 10 dinor-24-oxaoleana-20(29)-en-3-one in 2 mL of dry tetrahydrofuran is
 added and the solution is heated under nitrogen at 55°C for 14 h. The
 mixture is applied to a 10 cm column of silica gel, which is washed with
 2:1 ethyl acetate-hexane. The eluate is concentrated and purified by
 silica gel chromatography with 2:1 ethyl acetate-hexane to produce the
 15 title compound.


EXAMPLE 10

4-(2-Bromobenzoyl)oxy-6,7,15,16-tetrakis(acetyloxy)-18-hydroxy-22-
 20 methoxycarbonyl[6 α ,7 α ,15 β ,16 β]D:A-Friedo-A-homo-27,30-dinor-24-
oxaoleana-20(29),21-dien-3-one

- 66 -



Step A: 6,7,15,16-Tetrakis(acetyloxy)-21,22-epoxy-4, 18-dihydroxy-22-methoxycarbonyl[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]-D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-20(29)-en-3-one




A solution of 102.1 mg (0.130 mmole) of 4,6,7,15,16-pentakis(acetyloxy)-21,22-epoxy-18-hydroxy-22-methoxycarbonyl[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-20(29)-en-3-one in 4 mL of tetrahydrofuran and 2 mL of 3M aqueous HCl is heated at 40°C for 24h. The solution is diluted with dichloromethane and the layers are separated. The organic layer is washed with 0.1M phosphate buffer (pH 7), then is dried over MgSO₄

- 67 -

and concentrated. The residue is purified by silica gel chromatography with 2:1 ethyl acetate-hexane to produce the title compound.

5 **Step B:** 4-(2-Bromobenzoyl)oxy-6,7,15,16-tetrakis(acetyloxy)-
21,22-epoxy-18-hydroxy-22-methoxycarbonyl-
[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-
dinor-24-oxaoleana-20(29)-en-3-one



10


To a solution of 17.5 mg (23.5 μ mole) of 6, 7, 15, 16-tetrakis(acetyloxy)-21,22-epoxy-4, 18-dihydroxy-22-methoxycarbonyl-[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-20(29)-en-3-one in 0.5 mL pyridine is added 27.5 mL (237 μ mole) of benzoyl chloride. The solution is stirred at room temperature for 4 h, then is concentrated under reduced pressure. The residue is first filtered through a plug of silica gel and then purified by HPLC (Waters RCM, μ Porosil, 10 mm X 10 cm) using a mixture of 9.6:6 (5:4:1 hexane-methyl *tert*-butyl ether-acetonitrile:hexane) to produce the title compound.

15

20

- 68 -

Step C: 4-(2-Bromobenzoyl)oxy-6,7,15,16-tetrakis(acetyloxy)-18-hydroxy-22-methoxycarbonyl[6 α ,7 α ,15 β ,16 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-20(29),21-dien-3-one



5


A solution of 218 mg (0.55 mmole) of tungsten hexachloride in 8 mL of dry tetrahydrofuran is cooled to -78°C under nitrogen. Then 0.69 mL (1.10 mmole) of 1.6M butyllithium is added and the solution is allowed to warm to room temperature over 30 min. Then a solution of 50.2 mg (0.141 mmole) of 4-(2-bromobenzoyl)oxy-6,7,15,16-pentakis(acetyloxy)-21,22-epoxy-18-hydroxy-22-methoxycarbonyl-[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-20(29)-en-3-one in 2 mL of dry tetrahydrofuran is added and the solution is heated under nitrogen at 55°C for 14 h. The mixture is applied to a 10 cm column of silica gel, which is washed with 2:1 ethyl acetate-hexane. The eluate is concentrated and purified by silica gel chromatography with 2:1 ethyl acetate-hexane to produce the title compound.

20

- 69 -

WHAT IS CLAIMED IS:

1. A compound of structural Formula I:



5

I

or a pharmaceutically acceptable salt, crystal form or hydrate, wherein:

X is: O, S, NH or H and R¹;10 a is: a single bond, or a double bond when R⁴ is absent;

b and c are independently: a single bond or a double bond;

n is: 1 to 4;

15

m is: 1 to 4;

r is: 0 or 1;

20 s is: 0 or 1;

R¹ and R² are independently:

a) H, or

b) (C₁-C₆)-alkyl, wherein alkyl is unsubstituted or substituted
25 with one, two or three substituents selected from the group

- 70 -

- consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, vinyl, cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, aryl, wherein aryl is defined as phenyl or naphthyl,
- 5 unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, phenyl, phenoxy, cyano, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl and any two of
- 10 adjacent substituents can be joined to form a 5-, 6- or 7-membered fused ring said ring containing 1 or 2 oxygen atoms and the remainder carbon atoms, heteroaryl, wherein heteroaryl is defined as a 5 or 6-membered ring substituted with one and two heteroatoms selected from O, S, N,
- 15 unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, cyano, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, any two adjacent substituents can be
- 20 joined to form a 5-, 6- or 7-membered fused ring said ring containing 1 or 2 oxygen atoms and the remainder carbon atoms, or any two adjacent substituents can be joined together to form a benzo-fused ring;
- 25 R³ is:
- a) -(C₁-C₆)-alkyl, alkyl as defined above;
- b) -(C₁-C₆)-alkenyl, wherein alkenyl is unsubstituted or substituted with one, two or three substituents selected from
- 30 from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, aryl as defined above, and heteroaryl as defined above;
- c) -(C₁-C₆)-alkynyl, wherein alkynyl is unsubstituted or substituted with one, two or three substituents selected from

- 71 -

the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, aryl as defined above, and heteroaryl as defined above,


- 5 d) -aryl, aryl as defined above, or
 e) -heteroaryl, heteroaryl as defined above;

R⁴ is:

- 10 a) absent and a is a double bond;
 b) -H,
 c) -OH,
 d) =O,
 e) -O[(C=O)O_r]_sC₁-C₁₀-alkyl, alkyl as defined above,
 f) -O[(C=O)O_r]_sC₂-C₁₀-alkenyl, as defined above,
 g) -O[(C=O)O_r]_sC₂-C₆-alkynyl, alkynyl as defined above,
 h) -O[(C=O)O_r]_s(C₃-C₇)-cycloalkyl,
 i) -O[(C=O)O_r]_saryl, aryl as defined above,
 j) -O[(C=O)O_r]_sheteroaryl, heteroaryl as defined above,
 k) -O(CH₂)_nO(CH₂)_mheteroaryl, heteroaryl as defined above,
 l) -O(CH₂)_nO(CH₂)_maryl, aryl as defined above,
 m) -OC(=O)NR¹R²,
 n) -OSO₂R³, or
 o) -NR¹R².

- 72 -

2. The compound of structural Formula I, as recited in Claim 1,



I

5 or a pharmaceutically acceptable salt, crystal form or hydrate, wherein:

X is: O, S, or NH;

a is: a single bond;

10

b and c are independently: a single bond or a double bond;

n is: 1 to 4;

15 m is: 1 to 4;

r is: 0 or 1;

s is: 0 or 1;

20

R¹ and R² are independently:

a) H, or

b) (C₁-C₆)-alkyl, wherein alkyl is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, vinyl, cyano,

25

- 73 -

5 oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl,
CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl,
aryl, wherein aryl is defined as phenyl or naphthyl,
unsubstituted or substituted with one, two or three
10 substituents selected from the group consisting of: Br, Cl,
F, I, (C₁-C₆)-alkoxy, phenyl, phenoxy, cyano, nitro,
hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl,
CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl and any two of
adjacent substituents can be joined to form a 5-, 6- or 7-
15 membered fused ring said ring containing 1 or 2 oxygen
atoms and the remainder carbon atoms, heteroaryl, wherein
heteroaryl is defined as a 5 or 6-membered ring substituted
with one and two heteroatoms selected from O, S, N,
unsubstituted or substituted with one, two or three
20 substituents selected from the group consisting of: Br, Cl,
F, I, (C₁-C₆)-alkoxy, cyano, nitro, hydroxy, CHO, CO₂H,
COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R²,
NR¹COC₁-C₆-alkyl, any two adjacent substituents can be
joined to form a 5-, 6- or 7-membered fused ring said ring
25 containing 1 or 2 oxygen atoms and the remainder carbon
atoms, or any two adjacent substituents can be joined
together to form a benzo-fused ring;

R³ is:

- 25 a) -(C₁-C₆)-alkyl, alkyl as defined above;
 b) -(C₁-C₆)-alkenyl, wherein alkenyl is unsubstituted or
substituted with one, two or three substituents selected from
from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy,
cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl,
30 CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl,
aryl as defined above, and heteroaryl as defined above;
 c) -(C₁-C₆)-alkynyl, wherein alkynyl is unsubstituted or
substituted with one, two or three substituents selected from
the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy,

- 74 -

cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, aryl as defined above, and heteroaryl as defined above,


- 5 d) -aryl, aryl as defined above, or
 e) -heteroaryl, heteroaryl as defined above;

R⁴ is:

- 10 a) absent and a is a double bond;
 b) -H,
 c) -OH,
 d) =O,
 e) -O[(C=O)O_r]_sC₁-C₁₀-alkyl, alkyl as defined above,
 f) -O[(C=O)O_r]_sC₂-C₁₀-alkenyl, as defined above,
 g) -O[(C=O)O_r]_sC₂-C₆-alkynyl, alkynyl as defined above,
15 h) -O[(C=O)O_r]_s(C₃-C₇)-cycloalkyl,
 i) -O[(C=O)O_r]_saryl, aryl as defined above,
 j) -O[(C=O)O_r]_sheteroaryl, heteroaryl as defined above,
 k) -O(CH₂)_nO(CH₂)_mheteroaryl, heteroaryl as defined above,
 l) -O(CH₂)_nO(CH₂)_maryl, aryl as defined above,
20 m) -OC(=O)NR¹R²,
 n) -OSO₂R³, or
 o) -NR¹R².

- 75 -

3. The compound of structural Formula I, as recited in Claim 2,



I

5 or a pharmaceutically acceptable salt, crystal form or hydrate, wherein:

X is: O;

a is: a single bond;

10

b and c are independently: a single bond or a double bond;

n is: 1 to 4;

15 m is: 1 to 4;

r is: 0 or 1;

s is: 0 or 1;

20

R¹ and R² are independently:

a) H, or

b) (C₁-C₆)-alkyl, wherein alkyl is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, vinyl, cyano,

25

- 76 -

oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl,
 CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl,
 aryl, wherein aryl is defined as phenyl or naphthyl,
 unsubstituted or substituted with one, two or three
 5 substituents selected from the group consisting of: Br, Cl,
 F, I, (C₁-C₆)-alkoxy, phenyl, phenoxy, cyano, nitro,
 hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl,
 CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl and any two of
 10 adjacent substituents can be joined to form a 5-, 6- or 7-
 membered fused ring said ring containing 1 or 2 oxygen
 atoms and the remainder carbon atoms, heteroaryl, wherein
 heteroaryl is defined as a 5 or 6-membered ring substituted
 with one and two heteroatoms selected from O, S, N,
 15 unsubstituted or substituted with one, two or three
 substituents selected from the group consisting of: Br, Cl,
 F, I, (C₁-C₆)-alkoxy, cyano, nitro, hydroxy, CHO, CO₂H,
 COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R²,
 NR¹COC₁-C₆-alkyl, any two adjacent substituents can be
 20 joined to form a 5-, 6- or 7-membered fused ring said ring
 containing 1 or 2 oxygen atoms and the remainder carbon
 atoms, or any two adjacent substituents can be joined
 together to form a benzo-fused ring;

R³ is:

- 25 a) -(C₁-C₆)-alkyl, alkyl as defined above,
 b) -aryl, aryl as defined above, or
 c) -heteroaryl, heteroaryl as defined above;

R⁴ is:

- 30 a) -O[(C=O)O_r]_sC₁-C₁₀-alkyl, alkyl as defined above,
 b) -O[(C=O)O_r]_s(C₃-C₇)-cycloalkyl,
 c) -O[(C=O)O_r]_saryl, aryl as defined above,
 d) -O[(C=O)O_r]_sheteroaryl, heteroaryl as defined above,
 e) -O(CH₂)_nO(CH₂)_mheteroaryl, heteroaryl as defined above,

- 77 -

- f) $-\text{O}(\text{CH}_2)_n\text{O}(\text{CH}_2)_m\text{aryl}$, aryl as defined above,
- g) $-\text{OC}(=\text{O})\text{NR}^1\text{R}^2$, or
- h) $-\text{OSO}_2\text{R}^3$.


5 4. The compound of structural Formula I, as recited in Claim 3, or a pharmaceutically acceptable salt, crystal form or hydrate, wherein:

R⁴ is:

- 10 a) $-\text{O}[(\text{C}=\text{O})\text{O}_T]_s\text{aryl}$, wherein aryl is defined as phenyl or naphthyl, unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, phenyl, phenoxy, cyano, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, 15 CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl and any two of adjacent substituents can be joined to form a 5-, 6- or 7-membered fused ring said ring containing 1 or 2 oxygen atoms and the remainder carbon atoms, or
- 20 b) $-\text{O}[(\text{C}=\text{O})\text{O}_T]_s\text{heteroaryl}$, wherein heteroaryl is defined as a 5 or 6-membered ring substituted with one and two heteroatoms selected from O, S, N, unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, 25 cyano, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, any two adjacent substituents can be joined to form a 5-, 6- or 7-membered fused ring said ring containing 1 or 2 oxygen atoms and the remainder carbon atoms, or any two 30 adjacent substituents can be joined together to form a benzo-fused ring.

- 78 -

5. The compound of structural Formula I, as recited in Claim 1,



I

5 or a pharmaceutically acceptable salt, crystal form or hydrate, wherein:

X is: H and R¹;

a is: a single bond;

10

b and c are independently: a single bond or a double bond;

n is: 1 to 4;

15 m is: 1 to 4;

r is: 0 or 1;

s is: 0 or 1;

20

R¹ and R² are independently:

a) H, or

b) (C₁-C₆)-alkyl, wherein alkyl is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, vinyl, cyano,

25

- 79 -

5 oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl,
CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl,
aryl, wherein aryl is defined as phenyl or naphthyl,
unsubstituted or substituted with one, two or three
10 substituents selected from the group consisting of: Br, Cl,
F, I, (C₁-C₆)-alkoxy, phenyl, phenoxy, cyano, nitro,
hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl,
CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl and any two of
adjacent substituents can be joined to form a 5-, 6- or 7-
15 membered fused ring said ring containing 1 or 2 oxygen
atoms and the remainder carbon atoms, heteroaryl, wherein
heteroaryl is defined as a 5 or 6-membered ring substituted
with one and two heteroatoms selected from O, S, N,
unsubstituted or substituted with one, two or three
20 substituents selected from the group consisting of: Br, Cl,
F, I, (C₁-C₆)-alkoxy, cyano, nitro, hydroxy, CHO, CO₂H,
COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R²,
NR¹COC₁-C₆-alkyl, any two adjacent substituents can be
joined to form a 5-, 6- or 7-membered fused ring said ring
25 containing 1 or 2 oxygen atoms and the remainder carbon
atoms, or any two adjacent substituents can be joined
together to form a benzo-fused ring;

R³ is:

- 25 a) -(C₁-C₆)-alkyl, alkyl as defined above;
 b) -(C₁-C₆)-alkenyl, wherein alkenyl is unsubstituted or
substituted with one, two or three substituents selected from
from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy,
cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl,
30 CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl,
aryl as defined above, and heteroaryl as defined above;
 c) -(C₁-C₆)-alkynyl, wherein alkynyl is unsubstituted or
substituted with one, two or three substituents selected from
the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy,

- 80 -

cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl,
CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl,
aryl as defined above, and heteroaryl as defined above,


- 5 d) -aryl, aryl as defined above, or
 e) -heteroaryl, heteroaryl as defined above;

R⁴ is:

- 10 a) absent and a is a double bond;
 b) -H,
 c) -OH,
 d) =O,
 e) -O[(C=O)O_r]_sC₁-C₁₀-alkyl, alkyl as defined above,
 f) -O[(C=O)O_r]_sC₂-C₁₀-alkenyl, as defined above,
 g) -O[(C=O)O_r]_sC₂-C₆-alkynyl, alkynyl as defined above,
15 h) -O[(C=O)O_r]_s(C₃-C₇)-cycloalkyl,
 i) -O[(C=O)O_r]_saryl, aryl as defined above,
 j) -O[(C=O)O_r]_sheteroaryl, heteroaryl as defined above,
 k) -O(CH₂)_nO(CH₂)_mheteroaryl, heteroaryl as defined above,
 l) -O(CH₂)_nO(CH₂)_maryl, aryl as defined above,
20 m) -OC(=O)NR¹R²,
 n) -OSO₂R³, or
 o) -NR¹R².

- 81 -

6. The compound of structural Formula I, as recited in Claim 5,



I

5 or a pharmaceutically acceptable salt, crystal form or hydrate, wherein:

X is: H and R¹;

a is: a single bond;

10

b and c are independently: a single bond or a double bond;

n is: 1 to 4;

15 m is: 1 to 4;

r is: 0 or 1;

s is: 0 or 1;

20

R¹ and R² are independently:

a) H, or

b) (C₁-C₆)-alkyl, wherein alkyl is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, vinyl, cyano,

25

- 82 -

oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, aryl, wherein aryl is defined as phenyl or naphthyl, unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, phenyl, phenoxy, cyano, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl and any two of adjacent substituents can be joined to form a 5-, 6- or 7-membered fused ring said ring containing 1 or 2 oxygen atoms and the remainder carbon atoms, heteroaryl, wherein heteroaryl is defined as a 5 or 6-membered ring substituted with one and two heteroatoms selected from O, S, N, unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, cyano, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, any two adjacent substituents can be joined to form a 5-, 6- or 7-membered fused ring said ring containing 1 or 2 oxygen atoms and the remainder carbon atoms, or any two adjacent substituents can be joined together to form a benzo-fused ring;

R³ is:

- a) -(C₁-C₆)-alkyl, alkyl as defined above;
- b) -(C₁-C₆)-alkenyl, wherein alkenyl is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy, cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, aryl as defined above, and heteroaryl as defined above;
- c) -(C₁-C₆)-alkynyl, wherein alkynyl is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: Br, Cl, F, I, (C₁-C₆)-alkoxy,

- 83 -

cyano, oxo, nitro, hydroxy, CHO, CO₂H, COC₁-C₆-alkyl, CO₂C₁-C₆-alkyl, CONR¹R², NR¹R², NR¹COC₁-C₆-alkyl, aryl as defined above, and heteroaryl as defined above,

- d) -aryl, aryl as defined above, or
 5 e) -heteroaryl, heteroaryl as defined above;

R⁴ is:

- a) -OH,
 b) -O[(C=O)O_r]_sC₁-C₁₀-alkyl, alkyl as defined above,
 10 c) -O[(C=O)O_r]_s(C₃-C₇)-cycloalkyl,
 d) -O[(C=O)O_r]_saryl, aryl as defined above,
 e) -O[(C=O)O_r]_sheteroaryl, heteroaryl as defined above,
 f) -O(CH₂)_nO(CH₂)_mheteroaryl, heteroaryl as defined above,
 g) -O(CH₂)_nO(CH₂)_maryl, aryl as defined above,
 15 h) -OC(=O)NR¹R², or
 i) -OSO₂R³.

7. A compound selected from the group consisting of:

20 4,6,7,15,16-pentakis(acetyloxy)-18-hydroxy-22-methoxycarbonyl-
 [6 α ,7 α ,15 β ,16 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-
 1,20(29),21-trien-3-one;

25 4-(2-bromobenzoyl)oxy-6,7,15,16-tetrakis(acetyloxy)-18-hydroxy-22-
 methoxycarbonyl[6 α ,7 α ,15 β ,16 β]D:A-Friedo-A-homo-27,30-dinor-24-
 oxaoleana-1,20(29),21-trien-3-one;

30 4,6,7,15,16-pentakis(acetyloxy)-18-hydroxy-22-methoxycarbonyl-
 [6 α ,7 α ,15 β ,16 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-
 1,20(29),21-triene;

35 4,6,7,15,16-pentakis(acetyloxy)-21,22-epoxy-18-hydroxy-22-
 methoxycarbonyl[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-
 dinor-24-oxaoleana-20(29)-en-3-one;

- 84 -

4,6,7,15,16-pentakis(acetyloxy)-18-hydroxy-22-methoxycarbonyl-
[6 α ,7 α ,15 β ,16 β]D:A-Friedo-A-homo-27,30-dinor-24-oxaoleana-
20(29),21-dien-3-one;

5 4-(2-bromobenzoyl)oxy-6,7,15,16-tetrakis(acetyloxy)-18-hydroxy-22-
methoxycarbonyl[6 α ,7 α ,15 β ,16 β]D:A-Friedo-A-homo-27,30-dinor-24-
oxaoleana-20(29),21-dien-3-one;

6,7,15,16-tetrakis(acetyloxy)-21,22-epoxy-4,18-dihydroxy-22-
10 methoxycarbonyl[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-homo-27,30-
dinor-24-oxaoleana-20(29)-en-3-one; and

4-(2-bromobenzoyl)oxy-6,7,15,16-tetrakis(acetyloxy)-21,22-epoxy-18-
hydroxy-22-methoxycarbonyl[6 α ,7 α ,15 β ,16 β ,21 β ,22 β]D:A-Friedo-A-
15 homo-27,30-dinor-24-oxaoleana-20(29)-en-3-one.

8. A method treating a condition in a mammal, the
treatment of which is effected or facilitated by K_v1.3 inhibition,
comprising the administration, in an amount that is effective at
20 inhibiting K_v1.3, of a compound of Formula I.

9. The method of treating a condition in a mammal the
treatment of which is effected or facilitated by K_v1.3 inhibition, as
recited in Claim 8, wherein the condition is selected from the group
25 consisting of: resistance by transplantation of organs or tissue, graft-
versus-host diseases brought about by medulla ossium transplantation;
rheumatoid arthritis, systemic lupus erythematosus, Hashimoto's
thyroiditis, multiple sclerosis, myasthenia gravis, type I diabetes uveitis,
juvenile-onset or recent-onset diabetes mellitus, posterior uveitis,
30 allergic encephalomyelitis, glomerulonephritis, infectious diseases
caused by pathogenic microorganisms, inflammatory and
hyperproliferative skin diseases, psoriasis, atopic dermatitis, contact
dermatitis, eczematous dermatitis, seborrhoeic dermatitis, Lichen
planus, Pemphigus, bullous pemphigoid, Epidermolysis bullosa,

- 85 -

urticaria, angioedemas, vasculitides, erythemas, cutaneous eosinophilias, Lupus erythematosus, acne, Alopecia areata, keratoconjunctivitis, vernal conjunctivitis, uveitis associated with Behcet's disease, keratitis, herpetic keratitis, conical cornea, dystrophia epithelialis corneae, corneal leukoma, ocular pemphigus, Mooren's ulcer, Scleritis, Graves' ophthalmopathy, Vogt-Koyanagi-Harada syndrome, sarcoidosis, etc.; pollen allergies, reversible obstructive airway disease, bronchial asthma, allergic asthma, intrinsic asthma, extrinsic asthma and dust asthma, chronic or inveterate asthma, late asthma and airway hyper-responsiveness, bronchitis, gastric ulcers, vascular damage caused by ischemic diseases and thrombosis, ischemic bowel diseases, inflammatory bowel diseases, necrotizing enterocolitis, intestinal lesions associated with thermal burns and leukotriene B₄-mediated diseases, Coeliac diseases, proctitis, eosinophilic gastroenteritis, mastocytosis, Crohn's disease, ulcerative colitis, migraine, rhinitis, eczema, interstitial nephritis, Good-pasture's syndrome, hemolytic-uremic syndrome, diabetic nephropathy, multiple myositis, Guillain-Barre syndrome, Meniere's disease, polyneuritis, multiple neuritis, mononeuritis, radiculopathy, hyperthyroidism, Basedow's disease, pure red cell aplasia, aplastic anemia, hypoplastic anemia, idiopathic thrombocytopenic purpura, autoimmune hemolytic anemia, agranulocytosis, pernicious anemia, megaloblastic anemia, anerythroplasia, osteoporosis, sarcoidosis, fibroid lung, idiopathic interstitial pneumonia, dermatomyositis, leukoderma vulgaris, ichthyosis vulgaris, photoallergic sensitivity, cutaneous T cell lymphoma, arteriosclerosis, atherosclerosis, aortitis syndrome, polyarteritis nodosa, myocarditis, scleroderma, Wegener's granuloma, Sjogren's syndrome, adiposis, eosinophilic fascitis, lesions of gingiva, periodontium, alveolar bone, substantia ossea dentis, glomerulonephritis, male pattern alopecia or alopecia senilis by preventing epilation or providing hair germination and/or promoting hair generation and hair growth; muscular dystrophy; Pyoderma and Sezary's syndrome, Addison's disease, ischemia-reperfusion injury of organs which occurs upon preservation, transplantation or ischemic disease, for example, thrombosis and cardiac

- 86 -

infraction, endotoxin-shock, pseudomembranous colitis, colitis caused by drug or radiation, ischemic acute renal insufficiency, chronic renal insufficiency, toxins caused by lung-oxygen or drug, for example, paracort and bleomycins), lung cancer, pulmonary emphysema, cataracta, siderosis, retinitis, pigmentosa, senile macular degeneration, vitreal scarring, corneal alkali burn; dermatitis erythema multiforme, linear IgA bullous dermatitis and cement dermatitis, gingivitis, periodontitis, sepsis, pancreatitis, diseases caused by environmental pollution, aging, carcinogenesis, metastasis of carcinoma and hypobaropathy; disease caused by histamine or leukotriene-C₄ release; Behcet's disease, autoimmune hepatitis, primary biliary cirrhosis sclerosing cholangitis), partial liver resection, acute liver necrosis, necrosis caused by toxin, viral hepatitis, shock, or anoxia, B-virus hepatitis, non-A/non-B hepatitis, cirrhosis, alcoholic cirrhosis, hepatic failure, fulminant hepatic failure, late-onset hepatic failure, "acute-on-chronic" liver failure, augmentation of chemotherapeutic effect, preventing or treating activity of cytomegalovirus infection, HCMV infection, and antiinflammatory activity; and treatment of immunodepression or a disorder involving immunodepression, including AIDS, cancer, senile dementia, trauma, chronic bacterial infection, and certain central nervous system disorders.

10. The method as recited in Claim 9, wherein the condition is an autoimmune disease.

11. A method of preventing or treating the resistance to transplantation or transplantation rejection of organs or tissues into a patient in need thereof, which comprises the administration of a compound of Claim 1.

12. A method of suppressing the immune system in a subject in need thereof, which comprises the administration to the subject of an immune suppressing amount of a compound of Formula I, as recited in Claim 1.

- 87 -

13. A pharmaceutical formulation comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of the compound of Formula I, as recited in Claim 1 or a
5 pharmaceutically acceptable crystal form or hydrate thereof.

14. The pharmaceutical formulation of Claim 13, comprising in addition, a second immunosuppressive agent comprises azathioprine, brequinar sodium, deoxyspergualin, mizoribine,
10 mycophenolic acid morpholino ester, cyclosporin, FK-506 and rapamycin.

15. The method of Claim 12, comprising the coadministration of a second immunosuppressive agent.

15

16. A method of preventing or treating the resistance to transplantation or transplantation rejection of organs or tissues into a patient in need thereof, which comprises the administration of a compound of Claim 1.

20

17. A method of preventing or treating resistance by transplantation of organs or tissue, graft-versus-host diseases brought about by medulla ossium transplantation; rheumatoid arthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis,
25 myasthenia gravis, type I diabetes uveitis, juvenile-onset or recent-onset diabetes mellitus, posterior uveitis, allergic encephalomyelitis, glomerulonephritis, infectious diseases caused by pathogenic microorganisms, inflammatory and hyperproliferative skin diseases, psoriasis, atopic dermatitis, contact dermatitis, eczematous
30 dermatitises, seborrhoeis dermatitis, Lichen planus, Pemphigus, bullous pemphigoid, Epidermolysis bullosa, urticaria, angioedemas, vasculitides, erythemas, cutaneous eosinophilias, Lupus erythematosus, acne, Alopecia areata, keratoconjunctivitis, vernal conjunctivitis, uveitis associated with Behcet's disease, keratitis, herpetic keratitis, conical

- 88 -

- cornea, dystrophia epithelialis corneae, corneal leukoma, ocular pemphigus, Mooren's ulcer, Scleritis, Graves' ophthalmopathy, Vogt-Koyanagi-Harada syndrome, sarcoidosis, etc.; pollen allergies, reversible obstructive airway disease, bronchial asthma, allergic asthma,
- 5 intrinsic asthma, extrinsic asthma and dust asthma, chronic or inveterate asthma, late asthma and airway hyper-responsiveness, bronchitis, gastric ulcers, vascular damage caused by ischemic diseases and thrombosis. ischemic bowel diseases, inflammatory bowel diseases, necrotizing enterocolitis, intestinal lesions associated with thermal burns and
- 10 leukotriene B₄-mediated diseases, Coeliac diseases, proctitis, eosinophilic gastroenteritis, mastocytosis, Crohn's disease, ulcerative colitis, migraine, rhinitis, eczema, interstitial nephritis, Good-pasture's syndrome, hemolytic-uremic syndrome, diabetic nephropathy, multiple myositis, Guillain-Barre syndrome, Meniere's disease, polyneuritis,
- 15 multiple neuritis, mononeuritis, radiculopathy, hyperthyroidism, Basedow's disease, pure red cell aplasia, aplastic anemia, hypoplastic anemia, idiopathic thrombocytopenic purpura, autoimmune hemolytic anemia, agranulocytosis, pernicious anemia, megaloblastic anemia, anerythroplasia, osteoporosis, sarcoidosis, fibroid lung, idiopathic
- 20 interstitial pneumonia, dermatomyositis, leukoderma vulgaris, ichthyosis vulgaris, photoallergic sensitivity, cutaneous T cell lymphoma, arteriosclerosis, atherosclerosis, aortitis syndrome, polyarteritis nodosa, myocardosis, scleroderma, Wegener's granuloma, Sjogren's syndrome, adiposis, eosinophilic fascitis, lesions of gingiva, periodontium, alveolar
- 25 bone, substantia ossea dentis, glomerulonephritis, male pattern alopecia or alopecia senilis by preventing epilation or providing hair germination and/or promoting hair generation and hair growth; muscular dystrophy; Pyoderma and Sezary's syndrome, Addison's disease, ischemia-reperfusion injury of organs which occurs upon preservation,
- 30 transplantation or ischemic disease, for example, thrombosis and cardiac infraction, endotoxin-shock, pseudomembranous colitis, colitis caused by drug or radiation, ischemic acute renal insufficiency, chronic renal insufficiency, toxinsosis caused by lung-oxygen or drug, for example, paracort and bleomycins), lung cancer, pulmonary emphysema,

- 89 -

cataracta, siderosis, retinitis, pigmentosa, senile macular degeneration, vitreal scarring, corneal alkali burn; dermatitis erythema multiforme, linear IgA bullous dermatitis and cement dermatitis, gingivitis, periodontitis, sepsis, pancreatitis, diseases caused by environmental pollution, aging, carcinogenesis, metastasis of carcinoma and hypobaropathy; disease caused by histamine or leukotriene-C4 release; Behcet's disease, autoimmune hepatitis, primary biliary cirrhosis sclerosing cholangitis), partial liver resection, acute liver necrosis, necrosis caused by toxin, viral hepatitis, shock, or anoxia, B-virus hepatitis, non-A/non-B hepatitis, cirrhosis, alcoholic cirrhosis, hepatic failure, fulminant hepatic failure, late-onset hepatic failure, "acute-on-chronic" liver failure, augmentation of chemotherapeutic effect, preventing or treating activity of cytomegalovirus infection, HCMV infection, and antiinflammatory activity; and treatment of immunodepression or a disorder involving immunodepression, including AIDS, cancer, senile dementia, trauma, chronic bacterial infection, and certain central nervous system disorders which comprises the administration of a compound of Claim 1.

18. A method of treating a condition in a mammal, the treatment of which is effected or facilitated by K_v1.3 inhibition, comprising the administration of a pharmaceutical formulation comprising a pharmaceutical carrier and a compound of Formula I, in an amount that is effective at inhibiting K_v1.3.

19. A method of treating a condition in a mammal, the treatment of which is effected or facilitated by K_v1.3 inhibition, comprising the coadministration of a therapeutically effective amount of a compound of Formula I, as recited in Claim 1, with a second immunosuppressive agent.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/17481

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) :C07D 313/06; A61K 31/365 US CL :514/183; 549/266, 268 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 514/183; 549/266, 268 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAS		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US, A, 4,453,967 (MORI) 12 June 1984, see entire document.	1-19
Y	Phytochemistry, Vol. 29, No. 7, issued 1990, ABREU et al.; "A Nor-triterpenoid From Lophanthera Lactescens," pages 2257-2261, see pages 2257-2261.	1-19
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"A" document member of the same patent family	
"O" document referring to an oral disclosure, use, exhibition or other means		
"P" document published prior to the international filing date but later than the priority date claimed		
Date of the actual completion of the international search 17 JANUARY 1997	Date of mailing of the international search report 05 MAR 1997	
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer P.K. SRIPADA Telephone No. (703) 308-1235	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/17481

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/17481

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

- I. Claims 1-10, 13, 14, 18 and 19; drawn to compounds, pharmaceutical formulations and method of treating a condition in a mammal facilitated by K, 13 inhibitor.
- II. Claims 11, 16, 17 drawn to method of preventing or treating resistance of transplantation, classified in Class 514, subclass 183.
- III. Claims 12, 15 drawn to method of suppressing immune system; with coadministration of second agent, classified in Class 514, subclass 183.

The Groupings of the invention lack unity of invention because Groups I, II and III are drawn to different methods of use. If one of the methods is found in prior art, it would not make a reference to other methods, 37 CFR 1.475(d). The different methods of use have different etiologies, involving different cascade systems. The claims, therefore, are not so linked by a special technical feature within the meaning of PCT Rule 13.2 so as to form a single invention concept.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.